

TABLE 4
Regression line parameters

Pressure [kb]	<i>n</i>	<i>a</i> [km sec ⁻¹]	<i>b</i> $\frac{\text{km sec}^{-1}}{\text{g cm}^{-3}}$	$S_{(V,\rho)}$ [km sec ⁻¹]	<i>r</i>	<i>r</i> ² [%]
$V_p = a + b\rho$						
0.5	32	-3.55	3.32	0.20	0.96	92
1.0	32	-3.47	3.31	0.19	0.96	93
2.0	32	-3.35	3.28	0.18	0.97	94
6.0	32	-2.61	3.06	0.17	0.97	94
10.0	32	-2.01	2.88	0.17	0.96	92
$V_s = a + b\rho$						
0.5	32	-3.04	2.15	0.16	0.94	88
1.0	32	-2.73	2.06	0.15	0.94	89
2.0	32	-2.31	1.92	0.14	0.94	89
6.0	32	-1.53	1.66	0.14	0.92	85
10.0	32	-1.11	1.52	0.15	0.91	83
Pressure [kb]	<i>n</i>	<i>a</i> [g cm ⁻³]	<i>b</i> $\frac{\text{g cm}^{-3}}{\text{km sec}^{-1}}$	$S_{(V,\rho)}$ [g cm ⁻³]	<i>r</i>	<i>r</i> ² [%]
$\rho = a + bV_p$						
0.5	32	1.20	0.278	0.06	0.96	92
1.0	32	1.17	0.281	0.06	0.96	93
2.0	32	1.13	0.285	0.05	0.97	94
6.0	32	0.97	0.306	0.05	0.97	94
10.0	32	0.85	0.322	0.06	0.96	92
$\rho = a + bV_s$						
0.5	32	1.57	0.411	0.07	0.94	88
1.0	32	1.48	0.433	0.07	0.94	89
2.0	32	1.38	0.462	0.07	0.94	89
6.0	32	1.19	0.512	0.08	0.92	85
10.0	32	1.07	0.545	0.09	0.91	83

n = number of data points, $S_{(V, \rho)}$ = standard error of estimate of *V* on ρ , $S_{(\rho, V)}$ = standard error of estimate of ρ on *V*,*r* = correlation coefficient,*r*² = coefficient of determination.

In addition, the data has been filtered in accordance with the mean standard error of the refraction determinations. The distribution is similar to that found in earlier treatments [2, 9] and is clearly representative of the wide range in velocities observed for layer 2.

If the velocity histogram of fig. 5a is broken down into age groups as in figs. 5b and 5c, strikingly different velocity distributions can be seen for young and old sea floor. Young oceanic crust (fig. 5b) dis-

plays generally high layer 2 compressional wave velocities, the commonly observed values ranging from 4.5 to 6.3 km/sec with a mode of nearly 6.0 km/sec; old sea floor velocities (fig. 5c) are characteristically low, with common values ranging from 4.2 to 5.0 km/sec about a mode of 4.6 km/sec. Though it is possible to explain the entire spectrum of layer 2 velocities in terms of submarine weathering, it is evident from the wide range of velocities observed in each group that other processes such as neovolcanism, low grade metamorphism and fracturing are in part responsible for this range. Nonetheless, a pronounced trend of decreasing layer 2 refraction velocities with age similar to that observed in the laboratory is clearly indicated. We conclude that this trend, both in the laboratory and in the field, may be attributed to deep, progressive submarine weathering in oceanic layer 2 and that such weathering is consistent with the theory of sea floor spreading.

Acknowledgements

We wish to thank R. McConaghy and M. Mulcahey for their assistance with the measurements and for maintaining the high pressure equipment. This research was supported by National Science Foundation grant GA-36138. Samples were supplied through the assistance of the National Science Foundation.

References

- [1] N.I. Christensen and M.H. Salisbury, Sea floor spreading, progressive alteration of layer 2 basalts, and associated changes in seismic velocities, *Earth Planet. Sci. Letters* 15 (1972) 367.
- [2] R.W. Raitt, The crustal rocks, in: *The sea*, ed. M.N. Hill (Wiley, New York, 1963) 85.
- [3] D.A. McManus, R.E. Burns, O. Weser, T. Vallier, C.V. von der Borch, R.K. Olsson, R.M. Goll and E.D. Milow, Initial reports of the Deep Sea Drilling Project, Volume V (U.S. Government Printing Office, Washington, 1970).
- [4] A.G. Fisher, B.C. Heezen, R.E. Boyce, D. Bukry, R.G. Douglas, R.E. Garrison, S.A. Kling, V. Krashenninnikov, A.P. Lisitzin and A.C. Pimm, Initial reports of the Deep Sea Drilling Project, Volume VI (U.S. Government Printing Office, Washington, 1970).
- [5] E.L. Winterer, W.R. Riedel, P. Bronnimann, E.L. Gealy, G.R. Heath, L. Kroenke, E. Martini, R. Moberly, Jr., J. Resig and T. Worsley, Initial reports of the Deep Sea Drilling Project, Volume VII (U.S. Government Printing Office, Washington, 1971).

- [6] J.D. Hays, H.E. Cook III, D.G. Jenkins, F.M. Cook, J.T. Fuller, R.M. Goll, E.D. Milow and W.N. Orr, Initial reports of the Deep Sea Drilling Project, Volume IX (U.S. Government Printing Office, Washington, 1970).
- [7] F. Birch, The velocity of compressional waves in rocks to 10 kb, 1, *J. Geophys. Res.* 65 (1960) 1083.
- [8] G. Simmons, Velocity of shear waves in rocks to 10 kb, 1, *J. Geophys. Res.* 69 (1964) 1123.
- [9] G.G. Shor, Jr., H.W. Menard and R.W. Raitt, Structure of the Pacific basin, in: *The sea*, ed. A.E. Maxwell (Wiley, New York, 1971) 3.
- [10] N. Den, W.J. Ludwig, S. Murauchi, J.I. Ewing, H. Hotta, N.T. Edgar, T. Yashii, T. Asanuma, K. Hagiwara, T. Sato and S. Ando, Seismic refraction measurements in the northwest Pacific basin, *J. Geophys. Res.* 74 (1969) 1421.
J.I. Ewing, W.J. Ludwig, M. Ewing and S.L. Eittreim, Structure of the Scotia Sea and Falkland plateau, *J. Geophys. Res.* 76 (1971) 7118.
T.J.G. Francis and R.W. Raitt, Seismic refraction measurements in the southern Indian Ocean, *J. Geophys. Res.* 72 (1967) 3015.
T.J.G. Francis and G.G. Shor, Jr., Seismic refraction measurements in the northwest Indian Ocean, *J. Geophys. Res.* 71 (1966) 427.
- D.V. Helmberger and G.B. Morris, A travel time and amplitude interpretation of a marine refraction profile: primary waves, *J. Geophys. Res.* 74 (1969) 483.
- C.E. Keen and D.L. Barrett, A measurement of seismic anisotropy in the northeast Pacific, *Can. J. Earth Sciences* 8 (1971) 1956.
- X. Le Pichon, R.E. Houtz, C.L. Drake and J.E. Nafe, Crustal structure of the mid-ocean ridges, *J. Geophys. Res.* 70 (1965) 319.
- S. Murauchi, N. Den, S. Asano, H. Hotta, T. Yoshii, T. Asanuma, K. Hagiwara, K. Ichikawa, T. Sato, W.J. Ludwig, J.I. Ewing, N.T. Edgar and R.E. Houtz, Crustal structure of the Philippine Sea, *J. Geophys. Res.* 73 (1968) 3143.
- G.G. Shor, Jr., H.K. Kirk and H.W. Menard, Crustal structure of the Philippine Sea, *J. Geophys. Res.* 73 (1971) 2562.
- G.H. Sutton, G.L. Maynard and D.M. Hussong, Widespread occurrence of a high velocity basalt layer in the Pacific crust found with repetitive sources and sonobuoys, in: *The structure and physical properties of the earth's crust*, ed. J.G. Heacock, *Am. Geophys. Union Geophys. Monograph* 14 (1971) 193.