## B. CLEAVER AND D. E. MATHER

## COMPARISON OF EXPERIMENTAL $K_{\rm H}$ and $\Delta H$ values with those predicted by blander's theory

Blander <sup>2</sup> has used a theory originally proposed by Uhlig <sup>36</sup> to estimate the solubilities and heats of solution of inert gases in fluoride melts. The dissolution process is considered in three stages : (i) the gas is expanded from concentration  $C_g$  to concentration  $C_d$ . The free energy change is  $\Delta G_1 = \mathbf{R}T \ln (C_d/C_g)$ . (ii) The gas molecules are contracted to points and mixed with the liquid. The free energy change is  $\Delta G_2$ . (iii) The gas molecules are expanded to their proper size. The work required to create the cavities in the melt to accommodate the molecules is  $\Delta G_3 = 4\pi r^2 N\gamma$ , where *r* is the radius of the gas molecule and  $\gamma$  is the surface tension of the melt, here considered as a continuum.

Since the gas is in equilibrium with the liquid, the sum of these free energy changes is zero. If no chemical interaction occurs between the gas molecules and the ions in the melt,  $\Delta G_2$  may be neglected and

$$RT\ln\left(C_d/C_a\right) = -4\pi r^2 N\gamma. \tag{4}$$

Eqn (4) may be used to calculate the solubility of the gas at different temperatures, using experimental values of the surface tension. The heat of solution can be calculated from the temperature dependence of the solubility. This has been done for the systems studied here, and the results are included in table 1. The radii used in the calculations were <sup>2, 11</sup> (in units of  $10^{-10}$  m): He, 1.22; Ar, 1.92; N<sub>2</sub>, 2.00. This simple model predicts the solubilities to within an order of magnitude, and gives the heats of solution to within a factor of 2. The agreement is about as good as that obtained by Blander *et al.*,<sup>2</sup> and confirms the general validity of the model for inert gas solutions. The observed trends in our results are correctly predicted, i.e., (i) for a given gas and melt, the solubilities of a series of gases follow the inverse sequence of the molecular diameters. (ii) For a given gas dissolving in a series of melts, the solubilities increase with decreasing surface tension. This is exemplified by the series Ar+AgNO<sub>3</sub>, Ar+NaNO<sub>3</sub>, Ar+LiNO<sub>3</sub>, Ar+RbNO<sub>3</sub> (see table 1).

A feature of the results is that  $K_{\rm H}$  (expt.)/ $K_{\rm H}$  (calc.) is about 0.1 for He + NaNO<sub>3</sub> and for He + LiNO<sub>3</sub>. Blander <sup>2</sup> obtained a similarly low value for this ratio for He + NaF/ZrF<sub>4</sub>. As he suggests, this may reflect thermal motion of the gas atom in its cavity, so that the cavity radius is larger than the radius of the gas atom deduced from gas phase studies.

The broad agreement between theory and experiment increases the confidence with which the theory may be used to estimate gas solubilities in systems which have not been studied experimentally. Conversely, if experimental results differ greatly from the predicted values (e.g., if the solubility is more than a decade higher than predicted, or if  $\Delta H$  is negative), this may be taken as an indication that the free energy term  $\Delta G_2$  is not negligible.

## STANDARD ENTROPIES OF SOLUTION

Entropies of solution have been calculated from the experimental results for those systems which have been studied over a temperature range, and are listed in table 1.  $\Delta S^{\circ}$  is negative, and takes values between -8 and  $-24 \text{ J K}^{-1} \text{ mol}^{-1}$ . These are more negative than the corresponding entropy changes for inert gases dissolving in NaF/ZrF<sub>4</sub><sup>-1</sup> ( $\Delta S^{\circ} = 0$  to  $-6 \text{ J K}^{-1} \text{ mol}^{-1}$ ), for CO<sub>2</sub> in alkali halide melts <sup>5</sup> ( $\Delta S^{\circ} =$ -2.5 to  $-5 \text{ J K}^{-1} \text{ mol}^{-1}$ ), or for inert gases in benzene ( $\Delta S^{\circ} = -4$  to  $-7 \text{ J K}^{-1} \text{ mol}^{-1}$ ),

## 2482 SOLUBILITIES OF GASES IN MOLTEN NITRATES

but less negative than for inert gases in water <sup>38</sup> ( $\Delta S^{\circ} = -54 \,\mathrm{J}\,\mathrm{K}^{-1}\,\mathrm{mol}^{-1}$  for He, and  $-82 \text{ J K}^{-1} \text{ mol}^{-1}$  for argon, at 25°C). The large, negative values for aqueous solutions are generally attributed to structuring of the water surrounding the gas molecule, to give an ice-like cage. For nitrate melts, the fact that  $\Delta S^{\circ}$  is more negative than it is for halide solvents may possibly be due to some restriction of the rotational movement of the nitrate ions near the gas molecules.

We thank Dr. G. Saville for allowing us to use his unpublished PVT data for argon, and the Central Electricity Generating Board for an equipment grant and a Studentship (D. E. M.).

<sup>1</sup> W. R. Grimes, N. V. Smith and G. M. Watson, J. Phys. Chem., 1958, 62, 862.

<sup>2</sup> M. Blander, W. R. Grimes, N. V. Smith and G. M. Watson, J. Phys. Chem., 1959, 63, 1164.

3

<sup>3</sup> J. H. Shaffer, W. R. Grimes and G. M. Watson, J. Phys. Chem., 1959, 63, 1999.

<sup>4</sup> P. E. Field and J. H. Shaffer, J. Phys. Chem., 1967, 71, 3218.

<sup>5</sup> D. Bratland, K. Grjotheim, C. Krohn and K. Motzfeldt, Acta Chem. Scand, 1966, 20, 1811.

<sup>6</sup> H. von Wartenberg, Z. Elektrochem., 1926, 32, 330.

7 S. Allulli, J. Phys. Chem., 1969, 73, 1084.

<sup>8</sup> G. Bertozzi, EURATOM Report, no. EUR. 2495e, 1965.

<sup>9</sup> J. L. Copeland and W. C. Zybko, J. Phys. Chem., 1965, 69, 3631.
<sup>10</sup> J. L. Copeland and W. C. Zybko, J. Phys. Chem., 1966, 70, 181.

<sup>11</sup> J. L. Copeland and L. Seibles, J. Phys. Chem., 1966, 70, 1811.

<sup>12</sup> J. L. Copeland and S. Radak, J. Phys. Chem., 1967, 71, 4360.

<sup>13</sup> J. L. Copeland and L. Seibles, J. Phys. Chem., 1968, 72, 603.

14 J. L. Copeland and S. Radak, J. Phys. Chem., 1966, 70, 3356.

<sup>15</sup> J. L. Copeland and J. R. Christie, *J. Phys. Chem.*, 1969, 73, 1205.
<sup>16</sup> S. B. Tricklebank, L. Nanis and J. O'M Bockris, *Rev. Sci. Instr.*, 1964, 35, 807.

<sup>17</sup> M. K. Nagarajan and J. O'M Bockris, J. Phys. Chem., 1966, 70, 1854.

<sup>18</sup> A. F. M. Barton, B. Cleaver and G. J. Hills, Trans. Faraday Soc., 1968, 64, 208.

<sup>19</sup> G. J. Janz, Molten Salts Handbook (Academic Press, New York, London, 1967).

<sup>20</sup> P. E. Field, private communication ; J. Phys. Chem., in press.

<sup>21</sup> J. A. Beattie and O. C. Bridgeman, J. Amer. Chem. Soc., 1928, 50, 3133.

<sup>22</sup> J. L. Copeland, private communication.

<sup>23</sup> G. J. Janz, C. Solomons and H. J. Gardner, Chem. Rev., 1958, 58, 461.

<sup>24</sup> G. N. Lewis, and M. Randall, Thermodynamics, (McGraw Hill, New York, 2nd ed., 1961), p. 227. <sup>25</sup> I. R. Kritchevsky and J. S. Kasarnovsky, J. Amer. Chem. Soc., 1935, 57, 2168.

<sup>26</sup> G. Saville, private communication. The (*PV/RT*) values were computed by Dr. Saville from compressibility measurements by A. Lecocq, J. Rech. C.N.R.S., 1960, **11**, (no. 50), p. 55.

<sup>27</sup> E. P. Bartlett, results quoted as private communication in Int. Crit. Tables (McGraw Hill, New York, 1928), 3, 17.

<sup>28</sup> E. B. Smith and J. Walkley, J. Phys. Chem., 1962, 66, 597.

<sup>29</sup> Int. Crit. Tables, (McGraw Hill, New York, 1928), 3, 204.

<sup>30</sup> J. O. Hirschfelder, C. F. Curtiss and R. B. Bird, The Molecular Theory of Gases and Liquids, (John Wiley and Sons Inc., New York, 1954), p. 1110.

<sup>31</sup> B. Cleaver and B. C. J. Neil, Trans. Faraday Soc., 1969, 65, 2860.

<sup>32</sup> J. P. Frame, E. Rhodes and A. R. Ubbelohde, Trans. Faraday Soc., 1961, 42, 1075.

<sup>33</sup> A. N. Campbell and D. F. Williams, Can. J. Chem., 1964, 42, 1984.

34 L. C. Kenausis, E. C. Evers and C. A. Kraus, Proc. Nat. Acad. Sci., 1963, 49, 141.,

<sup>35</sup> G. Schlichthärle, Diss., (University of Karlsruhe, 1969), also G. Schlichthärle, K. Tödheide and E. U. Franck, J. Chim. Phys., Report. 19th Ann. Meeting. Soc. Chim. Phys. (Montpellier), 1968, p. 70.

<sup>36</sup> H. H. Uhlig, J. Phys. Chem., 1937, 41, 1215.

<sup>37</sup> R. P. Bell, Trans. Faraday Soc., 1937, 33, 496.

<sup>38</sup> D. D. Eley, *Trans. Faraday Soc.*, 1937, 35, 490.

PRINTED IN GREAT BRITAIN AT THE UNIVERSITY PRESS ABERDEEN