

phase change  
M.C.

H.J. Hall

CLOU EV61 0075

(Reprinted from *Nature*, Vol. 191, No. 4794, p. 1194 only,  
September 16, 1961)

### A New Phase formed by High-Pressure Treatment: Face-centred Cubic Molybdenum Monocarbide

The attempted preparation and retention of polymorphs dependent on pressure is an important area of high-pressure research. Some of these phases are found in Nature; others were unknown prior to the recent development of equipment suitable for research at high pressure and high temperature.

Although high-pressure phases are thermodynamically unstable at ambient conditions, a few have been retained by quenching to room temperature under pressure. High-pressure polymorphs found in the Earth's crust<sup>1</sup> include several minerals, for example, the pyroxene jadeite,  $\text{NaAlSi}_2\text{O}_6$  (ref. 2); kyanite,  $\text{Al}_2\text{SiO}_5$  (ref. 3); and diamond (ref. 4). Previously unknown phases are coesite<sup>2</sup>, a form of  $\text{SiO}_2$  with a higher density than quartz; borazon<sup>3</sup>, a cubic form of boron nitride; and several compounds of the garnet type<sup>1</sup>. A number of high-pressure polymorphs of pure metals, which unlike the above compounds cannot be retained at ambient conditions, have been reported<sup>7</sup>.

High-pressure experiments on the molybdenum-carbon system have revealed a new form for the monocarbide phase. The new form, designated  $\alpha\text{-MoC}$ , has been retained to atmospheric pressure. Synthesis was achieved from both an equiatomic mixture of molybdenum and carbon and an equiatomic mixture of the compound  $\text{Mo}_2\text{C}$  and carbon at pressures and temperatures in the range of 40–70 kilobars and 1,800°–2,500° C. The X-ray diffraction pattern for  $\alpha\text{-MoC}$  showed the first ten lines for a face-centred cubic structure. No additional lines were observed. The average lattice parameter was  $a_0 = 4.27 \text{ \AA}$ . In addition to high-pressure experiments in which the only product was  $\alpha\text{-MoC}$ , other experiments carried out at lower temperatures and pressures afforded evidence for all the phases reported<sup>8</sup> for the molybdenum-carbon system at atmospheric pressure. A summary of the lattice parameters of the various phases is given in Table 1.

Further experimental detail and a thermodynamic analysis of the effect of pressure on the equiatomic region of the molybdenum-carbon system will be published elsewhere.