

surface impedance with respect to magnetic field is plotted vs magnetic field in Fig. 3, which is a typical X-Y recording trace. The magnetic field for this run was oriented in the trigonal plane. We observe changes in the electron cyclotron mass of -4 to -8% /kilobar which can be compared with the results of Schirber.¹⁴ Schirber used deHaas-van Alphen studies in a high pressure environment to find a change of $\sim -6\%$ /kilobar for the smallest electron cyclotron mass in bismuth. Both results compare favorably within experimental error. More complete results are now being gathered for future publication.

* Indiana State University, Department of Physics, Terre Haute, Ind. 47809.

¹ C. S. E. Philips, J. Chem. Phys. **23**, 2388 (1955).

² T. Kushida, G. B. Benedek, and N. Bloembergen, Phys. Rev. **104**, 1364 (1956).

³ M. G. Vallauri and P. W. Forsbergh, Jr., Rev. Sci. Instrum. **28**, 198 (1957).

⁴ W. M. Walsh, Jr. and N. Bloembergen, Phys. Rev. **107**, 904 (1957).

⁵ A. W. Lawson and G. E. Smith, Rev. Sci. Instrum. **30**, 989 (1959).

⁶ I. P. Kaminow and R. V. Jones, Phys. Rev. **123**, 1122 (1961).

⁷ R. G. Goodrich, G. E. Everett, and A. W. Lawson, Rev. Sci. Instrum. **35**, 1596 (1964).

⁸ W. T. Doyle, T. F. Dutton, and H. P. Liu, Rev. Sci. Instrum. **40**, 171 (1969).

⁹ J. E. Schirber, *Physics of Solids at High Pressures* (Academic, New York, 1965), p. 46.

¹⁰ A. Okaya and L. F. Barash, IRE Proc. **50**, 2081 (1962).

¹¹ J. R. Clement and J. K. Logan, Advan. Cryog. Eng. **2**, 104 (1960).

¹² J. S. Dugdale, Nuovo Cimento **9**, 27 (1958).

¹³ A. F. Kip, D. N. Langenberg, and T. W. Moore, Phys. Rev. **124**, 359 (1961).

¹⁴ J. E. Schirber and W. J. O'Sullivan, Phys. Rev. B **2**, 2936 (1970)