

Table 1

The quantity $P_1 - P_2$ is the difference in calculated pressure at various temperatures, where

$$P_1 = 2.249[(T/T_0)^{1.528} - 1]$$

and

$$P_2 = 2.2293[(T/T_0)^{1.5351} - 1].$$

T (°K)	$P_1 - P_2$ (bar)
100	2.5
125	5.0
150	5.9
175	5.1
200	2.7
225	-1.4
250	-7.0
275	-14.4
300	-23.5
325	-34.1
350	-46.3
375	-60.2
400	-75.6
425	-92.7
450	-111.3
475	-131.4
500	-153.1
525	-176.3
550	-201.1
575	-227.3

Since Hardy, Crawford and Daniels' work has shown that the melting curve of mercury and argon cannot *both* be represented by a Simon melting equation, and since neither their fit nor the present one fits the data when extrapolated to

the 20-26 kilobar region (pending more and better data at these pressures), serious doubt has been cast upon the accuracy of the Simon equation's description of argon melting phenomena. Additionally, this raises some question as to the use of a Simon equation fitted to mercury melting data as a secondary pressure standard, a practice which has never been theoretically justified [10].

My thanks are extended to Dr. P. Bolsaitis and Dr. I. L. Spain for their advice; to the Center for Materials Research, University of Maryland, for a research stipend; to the University of Maryland Computer Science Center for the use of their facilities.

References

- [1] R. K. Crawford and W. B. Daniels, *J. Chem. Phys.* 50 (1969) 3171.
- [2] J. D. Grace and G. C. Kennedy, *J. Phys. Chem. Solids* 28 (1967) 977.
- [3] W. H. Hardy, R. K. Crawford and W. B. Daniels, *J. Chem. Phys.* 54 (1971) 1005.
- [4] A. van Itterbeek, O. Verbeke and K. Staes, *Physica* 29 (1963) 742.
- [5] P. H. Lahr and W. G. Eversole, *J. Chem. Eng. Data* 7 (1962) 42.
- [6] A. Michels, J. M. Levelt and W. deGraaf, *Physica* 24 (1958) 659.
- [7] A. Michels and C. Prins, *Physica* 28 (1962) 101.
- [8] D. W. Robinson, *Proc. Roy. Soc. A* 225 (1954) 393.
- [9] W. van Witzenburg and J. C. Stryland, *Can. J. Phys.* 46 (1968) 811.
- [10] E. C. Lloyd, C. W. Bockett and F. R. Boyd, *Science* 164 (1969) 860.

VIBRAT

Phil

An ir
by CO₂-
of the 0

The role
is of interest
lower laser
faster by ad
this letter m
of the lower
tures are re
relaxation r
laser power

A d.c. op
circuited wi
the laser tu
volt. The m
a few μ sec f
the gas part
nearly the g
used here no
level occurs
prevents the
[4]. The up
cited direct
relaxation o
and H₂O [3]
can be negl
through stir
relaxation c
rate equatio
it can be sh
cumstances
nential and
ation time c

The rela
been measu
for mixture
and 1-8.5 to
laser tube l
length and l
quartz tube
vapour from