

plane before and after rotation (selected so that $\beta > \alpha$); γ is the angle between the glide line and the axis of internal rotation and S_0 is the coefficient of resolved shear stress. For the glide system under consideration γ is 60° and S_0 is given values of 0.4-0.5 (Turner, et al., 1954). For rotations of L_0 ranging from 5° to 12° the calculated strains are 5 to 18 percent.

There are certain limitations to the acceptance of these figures as a measure of the actual strain in the rock. Estimation of strain from the rotation of L_0 lamellae assumes that all the deformation was achieved by translation-gliding on $\{0001\}$, and there is evidence in most grains of limited twin-gliding. Moreover, many of the grains contain minute dark granules which may have resulted from the disruption of rotated lamellae. It is probable, in the writer's opinion, that L_0 lamellae become too diffuse for measurement and are eventually disrupted when the strain is much in excess of the values recorded above. Thus the strain recorded in the rotation of L_0 lamellae in individual grains is probably less, and perhaps considerably less, than the total post-crystalline strain in the rock.

CONCLUSION

The Loch Ailsh dolomite has a strongly-oriented tectonite fabric of unknown origin, with almost orthorhombic symmetry. The fabric yields a unified picture of translation-gliding on $\{0001\}$ and twin-gliding on $\{02\bar{2}1\}$, dating from the final stage of deformation of the rock. This late post-crystalline deformation probably took place at a comparatively low temperature and was produced by a strong compression directed along an axis plunging 50° to $N315^\circ E$.

ACKNOWLEDGMENTS

The author is deeply grateful to Professor Arthur Holmes, for valuable advice during the course of the work, and to Professor F. J. Turner and Dr. G. Oertel, for constructive criticism of the manuscript. Thanks are also due to Dr. D. B. McIntyre, Dr. L. E. Weiss and Dr. P. C. Gilmour, for helpful discussion and advice. The work was carried out at the University of Edinburgh, with generous financial assistance from the University Court, the Carnegie Trust for the Universities of Scotland and the Cross Trust.

REFERENCES

- Borg, I., and Turner, F. J., 1953, Deformation of Yule Marble, Part 6: *Geol. Soc. America Bull.*, v. 64, p. 1343-1351.
- Bradley, W. F., Burst, J. F., and Graf, D. L., 1953, Crystal chemistry and differential thermal effects of dolomite: *Am. Mineralogist*, v. 38, p. 207-217.
- Christie, J. M., 1956, A tectonic study of the post-Cambrian thrusts of the Assynt region: Thesis, University of Edinburgh.
- Fairbairn, H. W., and Hawkes, H. E., Jr., 1941, Dolomite orientation in deformed rocks: *AM. JOUR. SCI.*, v. 239, p. 617-632.
- Felkel, E., 1929, Gefügestudien an Kalktektoniten: *Jahrb. d. Geol. Bundesanstalt*, v. 79, p. 33-85.
- Handin, J., and Fairbairn, H. W., 1955, Experimental deformation of Hasmark dolomite: *Geol. Soc. America Bull.*, v. 66, p. 1257-1273.
- Higgs, D. V., and Handin, J., 1954, Experimental deformation of dolomite single crystals (Abstract): *Geol. Soc. America Bull.*, v. 65, p. 1263.

- Johnsen, A., 1902, Biegungen und Translationen: Neues Jahrb. f. Min., Geol. Pal., v. 2, p. 133-153.
- Ladurner, J., 1953, Allgemeine Kennzeichnung und regionale Stellung alpinen Dolomittektonite: Jahrb. d. Geol. Bundesanstalt, v. 96, p. 253-300.
- Turner, F. J., 1953, Nature and dynamic interpretation of deformation lamellae in calcite of three marbles: AM. JOUR. SCI., v. 251, p. 276-298.
- Turner, F. J., Griggs, D. T., and Heard, H., 1954, Experimental deformation of calcite crystals: Geol. Soc. America Bull., v. 65, p. 883-934.
- Turner, F. J., Griggs, D. T., Heard, H., and Weiss, L. E., 1954, Plastic deformation of dolomite rock at 380°C: AM. JOUR. SCI., v. 252, p. 477-488.
- Turner, F. J., Griggs, D. T., Clark, R. H., and Dixon, H. R., 1956, Deformation of Yule Marble, Part 7: Geol. Soc. America Bull., v. 67, p. 1259-1294.

DEPARTMENT OF GEOLOGY
POMONA COLLEGE
CLAREMONT, CALIFORNIA