

rotational degrees of freedom being accessible to diatomic molecules but not to spherically symmetric rare gas atoms.<sup>2</sup>

Copeland and Zybko have observed Henry's law constants for high-pressure solubilities of Ar and He in molten  $\text{NaNO}_3$  at  $642^\circ\text{K}$ .<sup>5</sup> For Ar,  $K_{h,\text{Ar}} = (17.2 \pm 1.7) \times 10^{-7}$  mole  $\text{cm}^{-3}$  atm<sup>-1</sup>, and for He,  $K_{h,\text{He}} = (22.7 \pm 0.7) \times 10^{-7}$  mole  $\text{cm}^{-3}$  atm<sup>-1</sup>. The close-packed "spherical" radius of the  $\text{N}_2$  molecule is very nearly the same as that of Ar ( $1.92 \text{ \AA}$ ). Thus, the cavity creation work done against the *comparatively small* surface tension of  $\text{NaNO}_3$  by either gas would be expected to be small and approximately the same for each. However, the polarizability of  $\text{N}_2$  is  $1.734 \times 10^{-24}$  cm<sup>3</sup> molecule<sup>-1</sup> as compared to  $1.626 \times 10^{-24}$  cm<sup>3</sup> molecule<sup>-1</sup> for Ar.<sup>13</sup> On this basis, one would expect slightly greater ion-induced dipole solvation effects for  $\text{N}_2$  than for Ar, which would tend to make the former slightly more soluble if all other effects are nearly equal. The polarizability of He is only about

$0.2036 \times 10^{-24}$  cm<sup>3</sup> molecule<sup>-1</sup>,<sup>13</sup> and its close-packed radius is about  $1.22 \text{ \AA}$ .<sup>2</sup> Thus, much less work against surface tension is required to create a hole for a He atom than for either  $\text{N}_2$  or Ar. However, solvation effects for He would be practically nonexistent. It would appear, therefore, that the slightly greater solubility of He might be primarily the result of the relative ease of introducing such a small atom into the liquid, in spite of the partially compensating solvation effects which exist for the larger solute molecules.

*Acknowledgments.* The authors gratefully acknowledge support of this work by the National Science Foundation, Grant GP-4274. This paper has been based on the thesis of Lawrence Seibles, which has been submitted to the Graduate School of Kansas State University in partial fulfillment of the requirements for the degree of Master of Science.

(13) E. A. Moelwyn-Hughes, "Physical Chemistry," 2nd ed, Pergamon Press, Oxford, 1961, p 383.