

of $(\partial e / \partial p)_v$ provides no guarantee that C_v and $(\partial p / \partial T)_v$ are also constant. In fact different sets of values of C_v and $(\partial p / \partial T)_v$ that satisfy the condition of constant $(\partial e / \partial p)_v$ can be associated with different (T-p-v) equations of state that will give different values of shock temperature.

It is concluded that the values of shock temperature calculated along the isentropes are more realistic than those that would be calculated with models based on less thermodynamic information.

ACKNOWLEDGMENTS

The authors are grateful to Physical Research Laboratory, Edgewood Arsenal, for supporting this work under Contract DA-18-035-AMC-122(A), and thank Miss B. Loo for performing the numerical integration and plotting the diagrams.

REFERENCES

1. M. H. Rice, R. G. McQueen, and J. M. Walsh, "Compression of Solids by Strong Shock Waves," Solid State Physics, Vol. 6 (Academic Press, Inc., 1958).
2. J. M. Walsh and R. H. Christian, Phys. Rev. 97 (1544), 1955.
3. M. H. Rice and J. M. Walsh, J. Chem. Phys. 26 (824), 1957.
4. M. Cowperthwaite, Am. J. Phys. 34 (1025), 1966.
5. J. W. Gibbs, The Scientific Papers of J. Willard Gibbs; Vol. 1, Thermodynamics, Dover Publications, New York, 1960.
6. Ya B. Zeldovich and A. S. Kompaneets, Theory of Detonation, Academic Press, New York & London, 1960, pp 9-10.
7. R. Courant and K. O. Friedrichs, Supersonic Flow and Shock Waves, Interscience Publishers, New York, 1948.
8. R. G. McQueen and S. P. Marsh, "Equation of State for 19 Metallic Elements and Shock Wave Measurements to 2 Megabars," J. Appl. Phys., 31, 7, 1253-1269, 1960.
9. M. Cowperthwaite and J. H. Blackburn, "Temperature of Shocked Liquids," Stanford Research Institute Special Technical Report No. 14, Contract DA-18-035-AMC-122(A), 1967.
10. H. J. McSkimin, J. Acoust. Soc. Am. 29, 1185, 1957.