

The fits for $f(v)$ and $g(v)$ define the (e-p-v) equation of state and allow calculation of Hugoniot curves centered on, and isentropes passing through points on the $p=0$ isobar in the (p-v) plane. The discontinuous change in slope of the $g(v)$ function at $v = 1.01316$ cc/g is manifest in the shape of these curves in the neighborhood of this volume.

V. CALCULATIONS

Construction of Hugoniot Curves and Isentropes

The Hugoniot curves centered on $p = 0$ at -20°C , 25°C , 158.5°C , and 256°C were calculated directly with Eq. 14. The isentropes passing through $p = 0$ at 25°C , 158.5°C , 256°C , and 296°C were constructed by integrating Eq. 15 numerically with a Runge-Kutta technique. The -20°C , 158.5°C , and 256°C Hugoniots and the 296°C isentrope are shown in Fig. 4. The 25°C Hugoniot and the 25°C and 296°C isentropes are shown in Fig. 5.

Calculation of Temperature

Equation 9 was used to calculate the temperature along the isentropes passing through 25°C and 296°C on the atmospheric isobar. The values of temperature along these isentropes are listed in Table II. Calculation of temperature at points where isentropes intersect the 25°C Hugoniot defines values of shock temperature along this Hugoniot curve. The point of intersection ($T_c = 522.1^{\circ}\text{C}$, $p_c = 58$ kbar, $v_c = 0.661$ cc/g) of the 296°C isentrope and the 25°C Hugoniot is the highest point on the 25°C Hugoniot where shock temperature can be calculated with the present data. The temperature on the 25°C Hugoniot below 58 kbar can be calculated with the isentropes lying to the left of the 296°C isentrope. The temperature where the 256°C isentrope intersects the Hugoniot is 456.7°C , and the temperature where the 158.5°C isentrope intersects the Hugoniot is 291.5°C .

It is not possible to calculate temperature on the 25°C Hugoniot above 58 kbar without making further assumptions. The temperature along the Hugoniot above 58 kbar was calculated with constant C_v rather than by extrapolating the low pressure data further. The equation for shock temperature² above a point (T_c, v_c) on a Hugoniot centered at ($p_0 = 0, v_0$),