

excited and its heat capacity equals the Boltzmann constant k , whence

$$\gamma_{\infty} = \frac{\sum_i \gamma_i}{3N}$$

the simple average of the mode gammas. At intermediate temperatures one sees that the mode gammas are weighted by their associated Einstein heat capacities at the observation temperature. Let us perform a qualitative analysis, estimating mode gammas from Fig. 2 of HARDY and KARO.

Compression data from Slater yields about 4 percent change in volume to the transition pressure. The average shift of the collected acoustic mode peaks is about 15 percent yielding an average acoustic mode gamma -4 . Similarly, the average optical mode gamma is about $+0.7$ yielding a high temperature limiting value of $\Gamma_G = -3.3$ decreasing algebraically with decreasing temperatures. Since in equation (1) B_T , V and C_v must be positive, the implication of the negative gamma is that the thermal expansion coefficient have negative values at all temperatures. The thermal expansion coefficient of RbI has been measured over a wide temperature range.⁽⁴⁻⁶⁾ Its value becomes negative at temperatures below 10°K but even at 5°K the value of γ estimated from the expansion data is about -0.2×10^{-4} , still less negative than value -4 suggested by the lattice dynamical calculations. (The low temperature limit of γ calculated from measurements of the pressure dependence of the elastic constants of RbI is positive.)

BARRON's suggestion⁽⁷⁾ that the thermal expansion of solids is a quantity which yields information about the microscopic interactions in the solid

is relevant here. In the present case, it seems that the requirement to account for the experimental values of the thermal expansion and to find a mode frequency which will vanish to permit spontaneous NaCl-CsCl transition will provide a challenge for the lattice dynamicists. We feel at present that the mode instability hypothesis probably is incorrect as the explanation of the spontaneous nature of the transformation on RbI.

This work was supported in part by the Army Research Office (Durham).

Princeton University
Princeton
New Jersey

W. B. DANIELS

Note added by author in proof—Since submission of this paper, direct measurements have been reported by SAUDERSON, *Phys. Rev. Lett.* **17**, 530 (1966), using inelastic neutron scattering techniques at high pressure, of the shift of the Ta [100] mode energy with compression over the pressure range to the transformation. A decrease of only about 13 percent was observed, which is consistent with the suggestion of the present paper.

REFERENCES

1. DANIELS W. B., *Bull. Am. Phys. Soc.* **7**, 237 (1962).
2. DANIELS W. B. and SMITH C. S., *Physics and Chemistry of High Pressures*, Society of Chemical Industry, London (1963).
3. HARDY J. R. and KARO A. M., *Lattice Dynamics* (Ed. R. F. WALLIS) Pergamon Press, Oxford (1965).
4. SCHUELE D. E. and SMITH C. S., *J. Phys. Chem. Solids* **25**, 801 (1964).
5. WHITE G. K., *Proc. R. Soc. A286*, 204 (1965).
6. SWENSON C. A. Private communication.
7. BARRON T. H. K., *Ann. Phys.* **1**, 77 (1957) and *Phil. Mag.* **46**, 720 (1955).

... 28, pp. 350-351.
... and the NaCl-CsCl
... transition in RbI
... (20 July 1966)
... (1-3) that the spontaneous
... from the NaCl to the CsCl
... occurs via a structural
... shifting of the frequency of
... lattice vibrational mode at
... zone face, at a pressure
... formation. This problem is
... extended by HARDY and
... indicate that, for appropri-
... interactions, as the solid is
... transition conditions the
... the lattice vibrational
... shifted to lower energies
... are shifted slightly to
... in Fig. 2 of HARDY and
... of this note is to point out
... are strongly inconsistent
... (4-6) of thermal expans-
... ergy shifts calculated are
... range to the transition.
... quasi harmonic oscillator⁽⁷⁾
... excitations are considered
... nearly harmonic oscillators
... dependence of their
... This approximation leads
... isen's Gamma":

$$\gamma = \frac{\sum_i \gamma_i C_{vi}}{\sum_i C_{vi}}$$

expansion coefficient, B_T
 B_T/V is the heat capacity per

... ν_i
... V
... ν_i is any lattice vibra-
... C_{vi} is the Einstein heat
... de at the temperature of
... tion is over all oscillators
... vibrational spectrum. At
... oscillator is classically