

melting

H. J. Hall

DANI WB62 0082

PHYSICAL REVIEW LETTERS

VOLUME 8

JANUARY 1, 1962

NUMBER 1

LOW-TEMPERATURE LIMIT OF GRÜNEISEN'S GAMMA OF GERMANIUM AND SILICON

W. B. Daniels

Princeton University, Princeton, New Jersey

(Received November 28, 1961)

Gibbons¹ has measured the thermal expansion of Ge, Si, and InSb at low temperatures. Using these data and values of the heat capacity measured by various investigators, together with the Grüneisen relation,²

$$\gamma = \alpha V / \chi_T C_v, \quad (1)$$

he obtains a plot of γ vs T/θ_∞ reproduced as the solid lines of Fig. 1. α is the volume coefficient of thermal expansion, V the crystal volume, χ_T the isothermal compressibility, and C_v the heat capacity at constant volume. This plot indicates an anomalous negative peak for the case of Si and InSb, but not for Ge over the range of temperatures investigated. Gibbons has indicated by the dashed lines an extrapolation of γ toward zero as T/θ_∞ goes to zero. It is known that these materials with diamond-like structures reveal extraordinarily similar behavior in other lattice properties such as the temperature dependence of their Debye temperatures and even their lattice spectra.³ It seemed, then, worth investigating this situation wherein a difference of behavior was observed. Sheard⁴ discusses a way of obtaining high- and low-temperature limiting values of γ from a knowledge of the pressure dependence of the elastic constants of a solid, involving the averaging of a property over all the directions of a crystal. We have found a simple and quick method of obtaining the limiting value of γ as T approaches zero, from the following considerations. Derivation² of the relation (1) on the assumption that the Debye temperature is independent of temperature, which should be ex-

pected to be valid at very low temperatures, yields $-\gamma = d \ln \theta / d \ln V$. It is possible to calculate the limiting value of the Debye temperature at 0°K from the values of the elastic constants, molar volume, and density of a material; de Launay⁵ has prepared tables from which one can easily evaluate θ_0 using the relation:

$$\theta_0^3 = \frac{9N}{4\pi V} \left(\frac{h}{k} \right)^3 \left(\frac{C_{44}}{\rho} \right)^{3/2} \frac{9}{18 + \sqrt{3}} f(s, t), \quad (2)$$




FIG. 1. Variation of Grüneisen factor with reduced temperature T/θ_∞ for germanium ($\theta_\infty = 400^\circ\text{K}$), silicon ($\theta_\infty = 674^\circ\text{K}$), vitreous silica ($\theta_\infty = 495^\circ\text{K}$), and indium antimonide ($\theta_\infty = 214^\circ\text{K}$).¹