

the air bubble adhering to the lower part of one electrode shows that the liquid ahead of the shock is quite undisturbed. Plate 1, Figure 3, shows a shock wave which has travelled 10 mm into ethanol.

Plate 1, Figures 2 and 3, shows that the compressed liquid behind the shock front is transparent in both cases. This is an indication that freezing does not occur in spite of the high pressure in this region and is a confirmation of the finding of Walsh and Rice (1957) for water.

The actual pressure in the shock front cannot be determined directly but can be derived from the shock velocity and the free surface velocity in the shock. David and Hamann (1959) found in experiments in which similar explosive charges were used that the shock pressure at 5 mm distance from the end of the explosive was about 100,000 atm and the temperature 800 to 900° K. In those experiments the shock wave was however confined in a Polythene tube and since the shock propagation is very sensitive to the exact conditions surrounding the shock, these results are not directly applicable to the present experiments.

In the case of ethanol no shock velocity measurements under even approximately similar conditions are available for estimating the pressure. David and Hamann (1959, unpublished data) made some shock velocity measurements in methanol under the same conditions as in water and at 10 mm from the end of the explosive their results give a pressure of 64,000 atm in methanol compared with 68,000 in water. One would expect ethanol to give a similar pressure.

(b) Absorption Spectra

The photographic technique can also be used to investigate the absorption spectra of solutions under the extreme conditions obtained in shock waves. In order to do this the camera (11, see Fig. 1) is replaced by the entrance slit of the spectrograph and the lenses 8 are introduced in order to focus an image of the absorption cell on the slit.

The spectroscope used in the present work is a Hilger Constant Deviation Spectrograph which has an aperture of about $f/11$ and spreads the spectrum between 4000 and 6000 Å over 39 mm. Because of the strictly limited amount of light which can be obtained from an argon flash lasting 0.2 μ sec it was necessary to use a comparatively wide slit (0.05 mm) and very fast film (Ilford H.P.S.). Plate 2, Figures 1 and 2, shows the absorption spectra of 0.1M aqueous solutions of uranyl nitrate and neodymium chloride respectively. The four emission lines running across the spectrum are fortuitous and are due to the explosive flash other than that in the argon gap. They have been identified as the following lines: Na(D), 5890 Å (near red end of spectrum, not resolved); CaI, 4227 Å; CaII, 3968, 3933 Å (Petschek *et al.* 1955), and can serve as reference lines. In order to observe the solution both ahead of and behind the shock wave the full length (18 mm) of the spectroscope slit has to be used and this gives rise to the curved images on the photograph.

The lower part of Plate 2, Figure 1, shows the absorption bands in the blue which give uranyl nitrate solution its yellow colour. The upper portion shows the spectrum of the solution in the shocked condition. The absorption here is continuous

and extends into the green region of the spectrum. The pressure at the shock front in this photograph may be about 75,000 atm and the picture shows that a high pressure region extends behind the shock front for as far as is visible (5 mm) ahead of the opaque explosion products.

In Plate 2, Figure 2, the three equally spaced horizontal lines are due to 5 mm markers ruled on the front window of the absorption cell. The shock front is visible between the upper two markers. Near the bottom edge a mercury emission spectrum is superimposed. The wavelengths (in Å) marked along the top edge correspond to the maxima of the unperturbed absorption lines as determined on a microphotometer.

The absorption spectrum of an aqueous solution of neodymium chloride consists of a number of sharp lines and narrow bands and these are broadened and shifted to the red in the shocked solution. The pressure at the shock front in this case could be about 137,000 atm and the temperature about 1100°K. The temperature rise in the shock would alone account for the broadening of the absorption lines, but the red shift, amounting to between 20 and 30 cm^{-1} , can be attributed to the effect of pressure. This shift is very much smaller than those observed for transition metal complexes by Stephens and Drickamer (1959) at static pressures of up to 60,000 atm. The rare earth absorption which is due to excitation of well-embedded 4f electrons is presumably less affected by changes of environment caused by the high pressure than are the absorptions in the transition metals.

A detailed discussion of the spectroscopic results will be published later.

IV. ACKNOWLEDGMENTS

We wish to acknowledge the courtesy of the Department of the Army and the Chief Instructor of the School of Artillery, Sydney, who provided us with a location at which to carry out the experiments. We also wish to thank the School of Physics, University of Sydney, for the loan of the spectrometer.

V. REFERENCES

- COLE, R. H. (1948).—"Underwater Explosions." Ch. 6. (Princeton Univ. Press.)
COOK, M. A. (1958).—"The Science of High Explosives." Ch. 2. (Reinhold Publishing Corp.: New York.)
DAVID, H. G., and HAMANN, S. D. (1959).—*Trans. Faraday Soc.* **55**: 72.
DAVID, H. G., and HAMANN, S. D. (1960).—*Trans. Faraday Soc.* **56**: 1043.
PETSCHER, H. E., ROSE, P. M., GLICK, H. S., KANE, A., and KANTROWITZ, A. (1955).—*J. Appl. Phys.* **26**: 83.
STEPHENS, D. R., and DRICKAMER, H. E. (1952).—*J. Chem. Phys.* **30**: 1518.
SYKES, W. G. (1959).—*Science* **130**: 1051.
WALSH, J. M., and RICE, M. H. (1957).—*J. Chem. Phys.* **26**: 815.
WINNING, C. H., and EDGERTON, H. E. (1952).—*J. Soc. Mot. Pict. Telev. Engrs.* **59**: 178.