

nitrate illustrates how this technique may be used to study the pressure dependence of complex equilibria. Consider the complex equilibrium involving the formation of MX from ions M^+ and X^- . Electrostriction effects will in general lead one to conclude that the sum of the partial molar volumes of the ions will be less than the partial molar volume of the complex species [17]. Therefore, an increase in pressure will favour the dissociation of the complex MX. Conductivity studies tabulated by HORNE support this conclusion, e.g. the ratio of the dissociation constant at 1 kbar relative to the 0.001 kbar value for $FeCl_2^{2+}$ is 10, and for $MgSO_4$ is 1.5 [18]. The variation of spectral intensity of part of the Raman spectrum of thorium nitrate is shown in Fig. 3. The region between 1200 and 1600 cm^{-1} may be assigned to fundamental modes of vibration of solvated and complexed nitrate ions [19, 20]. The contour between 1200 and 1450 cm^{-1} , which may be ascribed to solvated nitrate ion, increases in intensity with increase in pressure. This effect may be contrasted with the decrease in intensity of the band at 1527 cm^{-1} , ascribed to complexed nitrate ion, with increase in pressure. These observations are interpreted in terms of the dissociation of complex nitrate-thorium species with increase of pressure. Consistent intensity variations are obtained for both rising and falling pressure.

Acknowledgements—The authors would like to acknowledge valuable discussion with Dr. J. A. KONINGSTEIN (Carleton University) and Dr. H. J. BERNSTEIN (N.R.C. Ottawa). Invaluable technical help was obtained from the Division of Physics Optical Shop, N.R.C., with regard to the sapphire windows, and from the Science Workshop of Carleton University, with regard to cell modification.

[17] R. A. HORNE, *Marine Chemistry*, p. 166. Wiley-Interscience, New York (1969).

[18] R. A. HORNE, *Advances in High Pressure Research* (Edited by R. S. BRADLEY), Vol. 2, p. 204. Academic Press, London (1969).

[19] D. E. IRISH and A. R. DAVIS, *Can. J. Chem.* **46**, 943 (1968).

[20] A. R. DAVIS and R. A. PLANE, *Inorg. Chem.* **7**, 2565 (1968).

The first part of the paper discusses the general theory of the subject, and the second part discusses the experimental results. The results show that the theory is in good agreement with the experimental data.

The authors are grateful to the following persons for their assistance in the preparation of this paper: Mr. A. B. C., Mr. D. E. F., and Mr. G. H. I.

Received for consideration, January 1, 1954.

The second part of the paper discusses the general theory of the subject, and the third part discusses the experimental results. The results show that the theory is in good agreement with the experimental data.

The authors are grateful to the following persons for their assistance in the preparation of this paper: Mr. A. B. C., Mr. D. E. F., and Mr. G. H. I.

Received for consideration, January 1, 1954.