Fig. 6. Isochrons (lines of equal time) for 99% conversion of Kamsdorf aragonite → calcite plotted in the calcite stability field. Isochron values are in minutes.

factor responsible for the difference in the gradient obtained by Brown et al. and by us appears to be the magnitude of the pressure effect on the transition. The former authors allow only a factor of 10 for a pressure effect 1 kb below the phase boundary.

At best such calculations are only approximate inasmuch as they rest on the assumption that the Franciscan rocks were dry during unloading and that the changing of pressure-temperature conditions of the aragonite material down the geothermal gradient (i.e., rate of unloading) was slow enough to allow sufficient time at the higher temperatures for transformation to take place.

Of further interest is the fact that the rates of the calcite → aragonite transformation at 375 to 405°C and 15 ± 1 kb are of the same order of magnitude as those for the reverse transformation at 400°C and 1 bar. It is apparent that the amount of overstepping of the equilibrium pressure has roughly the same effect in rate acceleration as the degree of understepping. One might also expect (although on a less certain experimental basis) that a set of isochrons similar to that proposed for the calcite field is applicable to the aragonite field.

REFERENCES

Ahrens, T. J., and S. Katz, Ultrasonic observation of the calcite-aragonite transition, *J. Geophys. Res., 68,* 529–537, 1963.

Bridgman, P. W., The Velocity of polymorphic changes between solids, *Proc. Am. Acad. Arts Sci., 52,* 57–88, 1915.

Brown, W. H., W. S. Fyfe, and F. J. Turner, Aragonite in California Glaucophane schists, and the kinetics of the aragonite-calcite transformation, *J. Petrol., 3,* 566–585, 1962.

Buerger, M. J., and E. Washken, Metamorphism of minerals, *Am. Mineralogist, 32,* 296–308, 1947.

Chaudron, G., Contribution a l'etude des reactions dans l'etat solide cinetique de la transformation aragonite-calcite, *Intern. Symp. Reactivity of Solids, Gothenburg,* 9–20, 1952.

Clark, S. P., A note on the calcite-aragonite equilibrium, *Am. Mineralogist, 42,* 564–566, 1957.

Davis, B. L., High-pressure X-ray investigation of $CaCO_3$–II and $CaCO_3$–III at 25°C, and of the calcite ⇌ aragonite transition in the 300–500°C temperature interval, Ph.D. dissertation, 133 pp., University of California, Los Angeles, 1964.

Davis, B. L., and L. H. Adams, X-ray diffraction evidence for a critical end point for cerium I and cerium II, *J. Phys. Chem. Solids, 25,* 379–388, 1964.

Griggs, D. T., M. S. Patterson, H. C. Heard, and F. J. Turner, Annealing recrystallization in calcite crystals and aggregates, in *Rock Deformation,* edited by D. T. Griggs and J. Handin, *Geol. Soc. Am. Mem., 79,* 382 pp., 1960a.

Griggs, D. T., F. J. Turner, and H. C. Heard, Deformation of rocks at 500°C to 800°C, in *Rock Deformation,* edited by D. T. Griggs and J. Handin, *Geol. Soc. Am. Mem., 79,* 382 pp., 1960b.

Jamieson, J. C., Phase equilibrium in the system calcite-aragonite, *J. Chem. Phys., 21,* 1385–1390, 1953.

Jamieson, J. C., Introductory studies of high-pressure polymorphism to 24,000 bars with some comments on calcite–II, *J. Geol., 65,* 334–343, 1957.

MacDonald, G. J. F., Experimental determination of calcite-aragonite equilibrium relations at elevated temperatures and pressures, *Am. Mineralogist, 41,* 744–756, 1956.