

neighbor Sn-Sn and Mg-Mg interactions. The requirements of symmetry greatly reduce the number of independent constants.

$$D^{\text{Sn core-Sn Shell}} = \begin{pmatrix} \delta & 0 & 0 \\ 0 & \delta & 0 \\ 0 & 0 & \delta \end{pmatrix}$$

$$D^{\text{Mg-Sn}} = \begin{pmatrix} \alpha_1 & \beta_1 & \beta_1 \\ \beta_1 & \alpha_1 & \beta_1 \\ \beta_1 & \beta_1 & \alpha_1 \end{pmatrix}$$

$$D^{\text{Sn-Sn}} = \begin{pmatrix} \alpha_2 & 0 & 0 \\ 0 & \beta_2 & \gamma_2 \\ 0 & \gamma_2 & \beta_2 \end{pmatrix},$$

$$D^{\text{Mg-Mg}} = \begin{pmatrix} \beta_3 & 0 & 0 \\ 0 & \beta_3 & 0 \\ 0 & 0 & \alpha_3 \end{pmatrix}.$$

Proc. 7th Int. Bull. Phys. Soc. Am. 30, 988 (1953).

Physics 7, 218
Semiconductors
Academic Press,
ELLER E., *Phys.*

Phys. Rev. 112, Proc. Int. Conf.
963).
Electronics 1, 629

HOSLER W. R.
109, 819 (1962).
Rev. 129, 2024

Soc. (London)
an. J. Phys. 40,
, 1955 (1949).
(1950).
K., *Phys. Rev.*

Lattice Dynamics,

²⁴⁾ we define the ion and another. lly as

) is evaluated at -body potential between one ion

ants for the Sn
Sn, next nearest