

16. Barral, II, E. M., Porter, R. S. and Johnson, J. F., *J. Phys. Chem.* **71**, 895 (1967).
17. Pincus, P., *J. de Physique (Paris)* **30**, C4, 8 (1969).
18. Maier, W. and Saupe, A., *Z. Naturforschg.* **14a**, 882 (1959); *Z. Naturforschg.* **15a**, 287 (1960).
19. Le Pesant, J. P. and Papon, P., these proceedings.
20. Blinc, R., O'Reilly, D. E. and Peterson, E. M., *Solid State Comm.* **6**, 839 (1968).
21. Bravo, N., Doane, J. W., Arora, S. L. and Fergason, J. L., *J. Chem. Phys.* **50**, 1398 (1969).
22. Jain, P. L., Lee, J. C. and Spence, R. D., *J. Chem. Phys.* **23**, 878 (1955); Rowel, J. C., Phillips, W. D., Melby, L. R. and Panar, M., *J. Chem. Phys.* **43**, 3442 (1965).
23. Theoretical estimates (see Ref. 26) of S_c for a hard rod gas yield $S_c \sim 0.85$.
24. Hoyer, W. A. and Nolle, A. W., *J. Chem. Phys.* **24**, 803 (1956).
25. μ_I can include short range order effects (but we neglect such effects altogether in a molecular field treatment).
26. Onsager, L., *Ann. N.Y. Acad. Sci.* **51**, 627 (1949); Zwanzig, R., *J. Chem. Phys.* **39**, 1714 (1963).
27. Chandrasekhar, S., Krishnamurti, D. and Madhusudana, N. V., *Mol. Cryst. and Liq. Cryst.* **8**, 45 (1969); Chandrasekhar, S., Shashidhar, R. and Tara, N., *Mol. Cryst. and Liq. Cryst.* **10**, 337 (1970).
28. The quadratic dependence in S is produced by a $\cos^2\theta$ angular dependence of the intermolecular anisotropic interactions (quadrupolar symmetry).⁽¹⁸⁾ Some evidence for the importance of higher powers in S with more "exotic" molecules has been given by P. G. James and G. R. Luckhurst, *Mol. Phys.* **19**, 489 (1970). In the case of hard rods, it is clear that all higher powers in S will be involved. On the other hand, terms linear in S would correspond to ferroelectricity or piezoelectricity in the nematic state, but these have not been observed.
29. Chow, L. C. and Martire, D. E., *J. Phys. Chem.* **73**, 1127 (1969).