

prove this by the fact that UH₃ has a higher Curie temperature¹⁵ and a higher magnetization¹⁶ than UD₃. The zero point motion is larger in UH₃ than in UD₃ and this results in an additional lattice expansion. Going back to Th₄H₁₅ we expect a small negative isotope effect in comparison to Th₄D₁₅. Unfortunately this cannot be detected because of difficulties in identically preparing the deuteride and hydride.

We believe our picture to be only a very rough one. No details are known, e.g. concerning the portion

of hydrogen electrons filling the thorium *f*-band and how this mechanism could be influenced by pressure.

In any case, additional measurements, especially of the position of the *f*-band relative to the Fermi level and of the electronic specific heat of Th₄H₁₅, would help to further clarify the situation.

Acknowledgements – We should like to thank Prof. W. Buckel for interesting discussions. One of us (C.B.S.) gratefully acknowledges research support from the Research Corporation and from the National Science Foundation.

REFERENCES

1. SATTERTHWAITE C.B. and TOEPKE I.L., *Phys. Rev. Lett.* **25**, 741 (1970).
2. BUCKEL W. and STRITZKER B., *Phys. Lett.* **43A**, 403 (1973).
3. BENNEMANN K.H. and GARLAND J.W., *Bull. Am. Phys. Soc.* **18**, 326 (1973).
4. SATTERTHWAITE C.B. and PETERSON D.T., *J. Less Common Metals* **26**, 361 (1972).
5. CATON R. and SATTERTHWAITE C.B., *Bull. Am. Phys. Soc.* **II19**, 348 (1974).
6. ZACHARIASEN W.H., *Acta Crystallogr.* **6**, 393 (1953).
7. JAYARAMAN A. *et al.*, *Rev. Sci. Instr.* **38**, 44 (1967).
8. CHU C.W., Dissertation, University of California, San Diego (1968).
9. SMITH T.F., *J. Low Temp. Phys.* **6**, 171 (1972).
10. DIETRICH M. (to be published).
11. SMITH T.F., CHU C.W. and MAPLE M.B., *Cryogenics* **9**, 53 (1969).
12. HILL H.H., *Nuclear Metallurgy*, Vol. 17 Part I, 1970 (Edited by MINER W.N.) Plutonium and other actinides p. 2.
13. KMETKO E.A. and HILL H.H., *Nuclear Metallurgy*, Vol. 17 Part I, 1970 (Edited by MINER W.N.) Plutonium and other actinides p. 233, and other contributions to this conference.
14. TRZEBIATOWSKI W., SLIWA A. and STALINSKI B., *Rocz. Chem.* **26**, 110 (1952); Further references are found in: LIBOWITZ G., *Metal Hydrides* p. 491. Academic Press, New York and London (1968).
15. FLOTOW H.E. *et al.*, *J. Amer. Chem. Soc.* **81**, 3529 (1959).
16. HENRY W.E., *Phys. Rev.* **109**, 1976 (1958).

Es wird über Messungen der Übergangstemperatur zur Supraleitung unter hydrostatischem Druck bis 28 kbar berichtet. Die lineare Anfangssteigung beträgt +42 mK/kbar.