

<sup>16</sup>G. E. Zil'berman, J. Exptl. Theoret. Phys. (U.S.S.R.) 21, 1209 (1951).

<sup>17</sup>G. E. Zil'berman and F. I. Itskovich, J. Exptl. Theoret. Phys. (U.S.S.R.) 32, 158 (1957), Soviet Phys. JETP 5, 119 (1957).

<sup>18</sup>S. Sidoriak and J. Robinson, Phys. Rev. 75, 118 (1949).

<sup>19</sup>L. Mackinson. Proc. Phys. Soc. B62, 170 (1949).

<sup>20</sup>T. Berlincourt and M. Steele, Phys. Rev. 95, 1421 (1954).

<sup>21</sup>F. Donahoe and F. Nix, Phys. Rev. 95, 1395 (1954).

<sup>22</sup>J. S. Dillon and D. Shoenberg, Phil. Trans. Roy. Soc. A248, 1 (1955).

Translated by P. F. Schmidt  
63

SOVIET PHYSICS JETP

VOLUME 35 (8), NUMBER 2

FEBRUARY, 1959

### ENERGY DEPENDENCE OF $\mu^+ - e^+$ DECAY ASYMMETRY

A. I. MUKHIN, E. B. OZEROV, and B. PONTECORVO

Joint Institute for Nuclear Research

Submitted to JETP editor March 3, 1958

J. Exptl. Theoret. Phys. (U.S.S.R.) 35, 340-347 (August, 1958)

The energy dependence of the asymmetry of electrons originating in the decay of polarized  $\mu$  mesons was studied with a set of scintillation counters biased against bremsstrahlung. The measurements were carried out for electron energies above 20 Mev. The observations are in a quantitative agreement (within a few percent) with the two-component neutrino theory. The experiments indicate that the degree of polarization of the  $\mu$  meson beam is  $0.81 \pm 0.11$ .

#### INTRODUCTION

THE hypothesis of parity nonconservation in weak interactions<sup>1</sup> has led<sup>2</sup> to a special variant of the neutrino theory, called the two-component or longitudinal neutrino theory. The attractiveness of such a theory makes it desirable to conduct experiments to decide whether the theory is correct or not.

A study of the spectrum of electrons produced in the decay of polarized  $\mu$  meson at various angles offers such an opportunity. It is, however, necessary to note that such an investigation can, in principle, only disprove the two-component neutrino theory, but cannot provide a final proof of its validity. In other words, a four-component neutrino theory can, for an appropriate choice of the coupling constants,<sup>3</sup> give a spectrum similar to that predicted by the two-component theory, namely

$$f(E, \theta) = 2E^2\{(3 - 2E) + \lambda(2E - 1) \cos \theta\}. \quad (1)$$

Here  $f(E, \theta)$  denotes the electron energy-distribution function,  $\theta$  is the angle between the elec-

tron momentum and the  $\mu$ -meson spin,  $E$  is the ratio of the electron energy to the maximum energy possible, and  $\lambda$  is a parameter that depends on the coupling constants.

Parity nonconservation in  $\pi - \mu - e$  decay was discovered in the classical experiments of Garwin, Lederman, and Weinrich<sup>4</sup> who showed, using electronic methods of particle detection, that  $\mu$ -meson beams from an internal target of a synchrocyclotron are polarized to a large degree, and that the angular distribution of electrons in  $\mu - e$  decay is of the form  $1 + a \cos \theta$ . It follows from the above that the two-component neutrino theory predicts a definite dependence of the factor  $a$  on the electron energy.

The energy dependence of the factor  $a$  found in reference 4 is weaker than that given by Eq. (1). The energy of the electrons was estimated in these experiments by determining the penetrating power of electrons passing through a graphite absorber. Taking it into account that the arrangement of counters in reference 4 was such that the bremsstrahlung radiation of the electrons could be re-