

FIG. 6. Dependence of pressure and type of electrolyte on cell emf at 150°C.
O, cell with 4.5 M CsCl; ●, cell with 4.5 M KCl.

1.43 kcal/mol respectively, whilst ΔV for these three temperatures (also at 1 Kbar) is -2.0_9 , -2.2_3 , -1.0_1 cm³. dE/dT 0.3 mV/deg and dE/dP 2.1₉, 2.3₅, 1.6₀ mV/Kbar, used for these calculations, were calculated from linear and quadratic fits of the relevant data given in Table 1.

From the cell-emf values given in Table 1, taking the values at the lowest molality of HCl it is possible to calculate the standard electrode potential of the skin-calomel electrode. This is done by simple subtraction, if the standard electrode potentials of the silver/silver-chloride electrode are known with sufficient precision. Lietzke, Greeley, Smith and Stoughton¹⁵ and recently Izaki *et al.*¹⁶ have made measurements to derive the standard electrode potential of the silver/silver-chloride electrode up to at least 200°C.

Table 2 shows the derived values of standard electrode potential of the calomel

TABLE 2. STANDARD ELECTRODE POTENTIALS OF THE SKIN-CALOMEL ELECTRODE FROM 25–200°C


Authors	$E^{\circ}, \text{Hg}_2\text{Cl}_2/\text{Hg}, \text{mV}$				
	25°C	70°C	100°C	150°C	200°C
Various ¹⁻¹⁹	267.93 to 268.23	—	—	—	—
This study (using Lietzke <i>et al.</i> ¹⁵)	268.0 ₄	249.5 ₄	232.9 ₄	195.9 ₄	144.3 ₄
This study (using Izaki <i>et al.</i> ¹⁶)	267.8 ₄	248.6 ₄	231.7 ₄	194.8 ₄	142.7 ₄

electrode over a range of temperature, from the present work. The standard electrode potential may be represented by

$E_{\text{Hg}_2\text{Cl}_2/\text{Hg}}^{\circ} = 276.261 - 0.30535T - 8.7170 \times 10^{-4}T^2 - 4.594 \times 10^{-6}T^3$ mV, with a scatter of ± 0.6 mV; this is the result of least-squares fit of the combined sets of results of Table 2, including the averaged value at 25°C from previous work.⁶⁻¹⁰ The rather large scatter is due to the large difference, ca 2 mV at 200°C, between the E° s of Lietzke *et al.*¹⁵ and those of Izaki.¹⁶ Selection of only one or other of the E° s would reduce this scatter, but no preference can really be made at the present time.

(d) Measurements involving the Orion chloride-reversible electrode

The Orion electrode (O.E.) was tested at 25 and 70°C and at 1–1900 bar. Table 3 shows values found for cells II and III at concentrations of 0.1 and 5 M HCl;



The potentials were measured simultaneously, at 1 s intervals; therefore, they should give the same potential as cell (I).

TABLE 3. EXPERIMENTAL CELL EMFS FOR CELLS (II) AND (III), USING THE CHLORIDE-REVERSIBLE ORION ELECTRODE

		Pressure, Kbar							
		0.001	0.050	0.50	1.00	1.50	1.70	1.90	
0.1 M HCl	25°C	E_{111}, mV	-5.1	—	+3.1	—	+2.65	-0.9	+3.8
		E_{111}, mV	-50.3	—	-42.4	—	-45.5	-49.9	-44.8
		$\Delta E = E_1$	+45.2	—	+45.5	—	+48.2	+49.0	+48.6
70°C		E_{111}, mV	—	-6.3	-6.3	-5.3	-5.1	-4.4	4.2
		E_{111}, mV	—	-67.0	-68.1	-67.8	-69.3	-68.1	-68.1
		$\Delta E = E_1$	—	+60.7	+61.8	+62.5	+64.2	+63.7	+63.9
5 M HCl	25°C	E_{111}, mV	-1.7	—	+2.7	+6.9	+10.5	+11.8	+13.2
		E_{111}, mV	+45.6	—	+42.2	+39.1	+36.3	+35.7	+34.7
		$\Delta E = E_1$	+43.9	—	+44.9	+46.0	+46.6	+47.5	+47.9
70°C		E_{111}, mV	+4.8	—	+2.5	+37.5	+51.5	+48.7	
		E_{111}, mV	+52.4	—	+56.4	+22.9	+10.3	+13.2	
		$\Delta E = E_1$	+57.2	—	+58.9	+60.4	+61.8	+61.9	

Thus, for example, taking the value of $E_{111} - E_{111}$ at 25°C, 0.1 M HCl and 0.5 kbar as 45.5 mV there is a 0.2 mV discrepancy with the value shown in Table 1. Similar discrepancies are apparent for other pressures and temperatures. However, the results in Table 3 demonstrate that any "asymmetry" potential the Orion electrode may possess is reasonably constant for short periods of time and that the electrode does function correctly as a chloride-reversible electrode under these conditions.

(e) Effect of nature of cation and concentration of electrolyte on cell potential

The Nernst equation for cell (I) does not depend on concentration or type of cation of the chloride electrolyte, as (I) shows. By reference to Table 1 it is seen that there is a variation in the cell emf for any particular concentration, temperature or pressure for the three different solutions studied. Thus the values given for 4.5 M HCl, KCl and CsCl at 100°C and 0.050 Kbar are 70.0, 71.1 and 69.3 mV; 76.0 and 73.6 mV were recorded for HCl at 100°C for 0.1 M HCl and 4.5 M HCl at 1.5 Kbar.

There are three possible reasons for this behaviour. The first may be disposed of quickly as being the most unlikely—that the anomalous results are a consequence of the presence of small amounts of bromide. Guntelburg¹⁷ warned of the dissimilar extent of sources of error imposed upon the silver/silver-chloride and classical calomel electrodes due to bromide impurity. In the present case precautions, already