2 at a edium y disfactor oils do higher

n oil, n the ensity 0°C is nay be nes by oressed ight in mpera-

75°

1.0190

1.0051
.9921
.9800
.9697
.9522
.9374
.9240
.9120
.8927

1.0227 1.0099 .9925 .9794 .9684 .9510 .9368

0 1.0178 4 1.0040 1 .9908 9 .9786 44 .9672 10 .9485 96 .9330 .9196




Fig. 1. Relative viscosity of lard oil as a function of volume.

ture and t_0 the time of fall at atmospheric pressure at 25°. The reader is referred to the original papers for details of method and computation.

DISCUSSION

It is apparent at once on inspection of Figs. 1, 2, and 3, which are drawn from the data of Table I, that the relative viscosity increases reatly with comparatively small decrease of volume. Inasmuch as the figures are drawn from data of different observers, as mentioned previously, there is some doubt as to the experimental accuracy of the viscosity-volume curves. However, the writer's acquaintance with the method sed by Kleinschmidt for the viscosity determinations leads him to estimate the inaccuracy the curves to be not more than a few percent, which does not seriously limit their applicability. The lack of serious deviation of the points repreenting the experimental values from the smooth arves of the figures gives an indication of the mobable degree of accuracy of the data.

An interesting feature of the figures is the rlative displacements of the viscosity-volume urves at 25°, 40° and 75°C. If viscosity were a function of volume only, the curves for each oil would coincide at all three temperatures. The fures show that this requirement is not satisfied any case. The viscosity-volume curve for lard at 25° departs from the curve at 75° by an




Fig. 2. Relative viscosity of sperm oil as a function of volume.




Fig. 3. Relative viscosity of Pennsylvania medium oil as a function of volume.

amount sufficient to change the viscosity by a factor of 2.3 at a volume of 0.99, and by a factor of 3.2 at a volume of 0.93. Similar curves for the Pennsylvania oil at 25° and 75° are even more relatively displaced; the discrepancy in viscosity varies from a factor of 3.8 at a volume of 0.99 to 7.6 at 0.94.