

HIGH-PRESSURE STUDIES OF THE SYSTEMS
 Mg_2GeO_4 - Mg_2SiO_4 AND GeO_2 - SiO_2 WITH SPECIAL
REFERENCE TO THE OLIVINE-SPINEL TRANSITION

(L. C. Card No. Mic 59-2880)

Frank Dachille, Ph.D.

The Pennsylvania State University, 1959

Phase equilibrium studies have been extended by more than one order of magnitude beyond pressures attainable in hydrothermal systems. Pressures close to 900,000 psi have been obtained by the use of relatively simple uniaxial devices in which a large force is applied to a sample for tens of hours by means of very hard tool steel or cemented carbide pistons with small bearing surfaces. The sample is pressed into a thin nickel ring and isolated from the piston faces by platinum foils. Temperatures of 550-600°C are the practical upper limits at the extreme pressure range. Tests have been made which show that the pressure is uniform across the sample assembly, and further, that the results are comparable to those obtained with hydrostatic pressure.

The system Mg_2GeO_4 - Mg_2SiO_4 above 500°C has been studied to the limits of the hydrothermal and uniaxial methods. It has been determined that the olivine-spinel phase transition is reversible and reproducible. The univariant p-t curve for this transition for Mg_2GeO_4 passes through 823°C at 10,000 psi and increases at the rate of $24.6 \pm 1.2^\circ C$ to 80,000 psi. The ΔV of the two phases at room temperature and pressure is 3.5cc per mol. The ΔH of the transition calculated via the Clapeyron relation is 3.7 Kcal/mol. The maximum silicate content of the spinel solid solution obtained is 53 mol percent at 65 kilobars and 542°C. Extrapolation of the phase fields shows that Mg_2SiO_4 should convert to the spinel form at 100 ± 15 kilobars at 542°C. Other considerations suggest that the transition temperatures should rise 13°C per kilobar.

Assuming linear variation of x-ray "d-spacing" data with composition for the spinel solid solutions, the spinel phase of Mg_2SiO_4 will have a cell edge of 8.22Å. Thus the ΔV of transition for Mg_2SiO_4 would be only 2.0 cc per mol.

The geophysical implication of these results is that an olivine-spinel transition in the mantle of the earth does appear to be a likely explanation of the seismic and density discontinuities at the 400 km depth as suggested by Jeffreys and Bernal provided that the mantle composition is in fact olivine, and that the temperatures in the mantle are in the range proposed by Daly or Turner and Verhoogen.

Uniaxial high pressure studies in the system GeO_2 - SiO_2 at 500°C locate the quartz-coesite transition for SiO_2 at 20.4 ± 0.4 kilobars. The quartz solid solution field is reduced from 33 mol percent GeO_2 at low pressures to es-

essentially nothing at the eutectoid pressure of 32 kilobars and 13 mol percent GeO_2 , leaving SiO_2 (coesite) and GeO_2 (rutile).

A beginning has been made in the use of infra-red data for diagnostic purposes in the solid state. An empirical relation has been worked out which allows comparison of coordination polyhedra about major cations in simple compounds. The infra-red data lead to the conclusion that the 4^+ cations of silicates and germanates with the olivine structure are situated in distorted tetrahedra characterized by two pairs of oxygens in different equivalent coordinations to the cation. It also leads to the proposal that the Mg_2SiO_4 spinel will have the normal structure, and that Mg_2GeO_4 spinel has the inverse structure. Some x-ray corroboration of this fact has been found in relative intensities of the (422) and (400) reflections. According to earlier calculations for Mg_2GeO_4 spinel, the type may be distinguished by the I422/I400 ratio and would be normal, disordered or inverse if the ratio were 1.67, 3.3 or 6.6 respectively. In the present work the ratio found was always above 3.8 and as high as 6.5 for Mg_2GeO_4 .

Microfilm copy of complete manuscript of 83 pages, available from University Microfilms, Ann Arbor, Michigan, \$2.00. Enlargements 6" x 8", 4¢ per page.