

REFERENCES

1. P. W. Bridgman, Proc. Am. Acad. Arts Sci. 72, 45 (1937).
2. C. W. F. T. Pistorius, J. B. Clark, and E. Rapoport, J. Chem Phys. 48, 5123 (1968).
3. H. Suga, T. Matsuo, and S. Seki, Bull. Chem. Soc. Japan 38, 1115 (1965).
4. D. L. Price, J. M. Rowe, J. J. Rush, E. Prince, D. G. Hinks, and S. Susman, J. Chem. Phys. 56, 3697 (1972). ✓
5. C. W. F. T. Pistorius, J. Phys. Chem. Solids 32, 2761 (1971).
6. P. W. Richter and C. W. F. T. Pistorius, Acta Cryst. B28, 3105 (1972).
7. M. Atoji, J. Chem. Phys. 54, 3514 (1971).
8. This is argued from an assumed R ln 8 configurational entropy in phase III and the change in entropy from slope and volume change along the III-IV boundary.
9. R. M. Brugger, R. B. Bennion, and T. G. Worlton, Phys. Lett. A24, 714 (1967).
10. G. Roult, R. Beyerlein, D. Decker, T. G. Worlton, Bull. of APS 18, 35 (1973).
11. D. G. Hinks, D. L. Price, J. M. Rowe and S. Susman, J. Crystal Growth 15, 227 (1972).
12. R. M. Brugger, R. B. Bennion, T. G. Worlton, W. R. Myers, Trans. Amer. Cryst. Ass. 5, 141 (1969).
13. D. L. Decker, J. Appl. Phys. 42, 3239 (1971).
14. D. L. Decker, and T. G. Worlton, J. Appl. Phys. 43, 4799 (1972).
15. The lattice parameters of Ge and Si were taken to be 5.65754⁰A and 5.4307 A respectively.