VOLUME 11, NUMBER 4

15 FEBRUARY 1975

Diffusion of copper in lead

D. L. Decker

Department of Physics and Astronomy, Brigham Young University, Provo, Utah 84602 (Received 29 October 1974)

This paper presents a reanalysis of experimental data on the diffusion of Cu in Pb. The new values of the activation energy and volume are 8.17 ± 0.11 kcal/mole and 0.16 ± 0.02 atomic volumes, respectively.

In 1972, Candland, Decker, and Vanfleet¹ published the measurement of the effects of pressure on the diffusion of Cu in Pb. They reported an activation energy of $\Delta H = 5.6$ kcal/mole at atmospheric pressure and suggested that Cu diffuses in Pb by a pure interstitial mechanism. Their value of the activation energy, however, is not in good agreement with atmospheric-pressure measurements of Dyson, Anthony, and Turnbull² and Miller.³ In attempting to correlate the measurement of the diffusion of the noble metals in Pb. using an extension of the theory of Miller, ⁴ we found⁵ that the value of 5.6 kcal/mole for ΔH of Cu diffusing in Pb was inconsistent with the data for the other noble metals, while the value of 8 kcal/mole given by Dyson et al.² was consistent. It seemed that the analysis of the high-pressure data to obtain ΔH must be in error, so I reexamined the original data of Candland et al.¹ It was immediately apparent that Candland et al.'s earlier atmospheric-pressure measurements were taken prior to their preanneal technique and showed definite non-Gaussian penetration profiles. If I removed these points from their work, their remaining atmospheric-pressure points agreed well with Miller and Dyson *et al*. The lowest-temperature points along each of Candland et al.'s isobars had diffusion values higher than the higher-temperature measurements. The corresponding diffusion profiles had less than one decade of penetration and so these points were rejected. Their remaining data points were analyzed along with the atmospheric-pressure points of Miller and Dyson *et al*. to yield the following results⁶: $\Delta H = 8.17 \pm 0.11$ kcal/mole, $D_0 = 0.0086 \pm 0.0009 \text{ cm}^2/\text{sec}, \Delta V/V_0$ = 0.16 ± 0.02, $\delta(\Delta V/V_0)/\delta P = -(2.9 \pm 1.7) \times 10^{-3}$ kbar⁻¹, and $\delta(\Delta V/V_0)/\delta T = (0.8 \pm 0.4) \times 10^{-4} \text{ K}^{-1}$. A graph of the data is shown in Fig. 1. Another surprising change in this analysis is the considerably larger value for $\Delta V/V_0$ than previously reported. If Candland et al. had realized that this value was so large, they might have been more hesitant in claiming that the diffusion of Cu in Pb was pure interstitial.

After submitting this paper an article⁷ was published which also considers Candland *et al.*'s original data. By rejecting data taken prior to the preanneal procedure, which was designed to avoid oxidation problems at the surface, Mundy *et al.*⁷ found a value of 7.0 ± 0.5 kcal/mole for the activation energy from Candland *et al.*'s data. Using only the four atmospheric-pressure data points which we included in this reanalysis, we would obtain $\Delta H=7.9\pm0.4$ kcal/mole, which agrees with the conclusions of Mundy *et al.* As they suggested, the earlier measurements of Candland *et al.* may have been distorted by solubility problems.




FIG. 1. Temperature dependence of the diffusion of Cu in Pb along several isobars. The solid lines are the computer fit to the data using Eq. (6) in Ref. 6. The open circles at atmospheric pressure are measurements of Ref. 1, while the solid data points come from Refs. 2 and 3. \Box , at 20.6 kbar; \triangle , at 27.0 kbar; \diamond , at 33.3 kbar; ∇ , at 40.1 kbar.

11

1770

¹C. T. Candland, D. L. Decker, and H. B. Vanfleet, Phys. Rev. B 5, 2085 (1972).

²B. F. Dyson, T. R. Anthony, and D. Turnbull, J. Appl. Phys. <u>37</u>, 2370 (1966).

³J. C. Miller, Argonne report (1972) (unpublished).

⁴J. W. Miller, Phys. Rev. <u>188</u>, 1074 (1969).

⁵D. L. Decker and H. B. Vanfleet, Bul. Am. Phys. Soc.

18, 428 (1973).

⁶J. A. Weyland, D. L. Decker, and H. B. Vanfleet, Phys. Rev. B 4, 4225 (1971).

⁷J. N. Mundy, J. W. Miller, and S. J. Rothman, Phys. Rev. B <u>10</u>, 2275 (1974). This is the same datum as in Ref. 3.

the aphynithe certify and whate of I'

Alle solution delayed in a paper on activity and outthe bed when also consider the continue of all is carly und then in a rejection data intention of the conanal 1 (rescalary, with a merical material to a roadoblightion produces, with a merical material to a roadbound a value of 7, 0 - 0 is attained in (or the perfration energy from Continued of the 2 data. To ing both the terms from Continued of the 2 data. To ing attain we included in the event where the terms and the second of the strate merical and the second the strate and the strate of the strate of the second the strate and the strate of the strate of the strate the strate and the strate of the strate of the strate of the data is a road of the strate of the strate of the strate of the data is a road of the strate of the strate of the strate of the data is a road of the strate of the strate of the strate of the the strate of the strate of the strate of the strate of the strate the strate of the stra



23. J. J. Trinjeradiré depension de châle diffusion of qui mitrichen en de 1 objets. The edit floor the unramatic difficient contract for the indef to dive transitie difficient de contract for the indef to dive tract de dire de contract de contract and the substantic direct oper, of the Product of the dist rest of the direct oper, of the Product of the dist. Tract of the direct.

11.13.31

11

DECK ~ DL 75- 0029