4860 atmospheres (pressure $= -(dE/dr_0)/4\pi r_0^2$). The shifts of spectral lines could also have an influence on the Lorentz–Lorenz function. In fact the behaviour of the Lorentz–Lorenz expression [8]) as function of density could be partially explained by assuming that the wave lengths of spectral lines diminish by an amount of some percents or some tens of percents (cf. also [9]) and [10])).

Our thans are due to Mr A. Botzen for valuable help with numerical calculations.

REFERENCES

1) A. Michels, J. de Boer and A. Bijl, Physica **4**, 981,1937 (Van der Waals-number).
2) A. Sommerfeld and H. Welker, Ann. Physik [5], **32**, 56, 1938.
3) E. T. Whittaker and G. N. Watson, Modern Analysis, page 337, Cambridge 1927.
4) H. Buchholz, Z. angew. Math. Mech. **23**, 47 and 101, 1943 (formula (5.5)).
5) E. Jahnke and F. Emde, Tables of Functions, page 1, 3rd edition, Leipzig-Berlin 1938.
6) H. A. Webb and J. R. Airey, Phil. Mag. **36**, 129, 1918.
7) Rep. Brit Ass. Advancement Sc. **94**, 276, 1926 (Oxford); **95**, 220,1927 (Leeds).
8) J. Hamers, thesis, Amsterdam 1941.
9) A. Michels and J. Hamers, Physica **4**, 995, 1937; A. Michels, H. and L. Lebesque and S. R. de Groot, Physica. in the press; A. Michels, A. Botzen and S. R. de Groot, Physica, in the press.
10) S. R. de Groot and C. A. ten Seldam, Physica 1947, letter to the editor.