

Advan. High Pressure Res. 3, 1-39  
(1969)

x	
CHAPTER 4	
Electrical Conductivity and Electronic Transitions at High Pressures	
N. H. MARCH	
I. Outline . . . . .	241
II. Pressure Ionization and Electronic Transitions . . . . .	244
III. Metallic Transitions at Lower Pressures . . . . .	254
IV. Electrical Conductivity of Alkali Metals . . . . .	263
V. Pressure-dependence of Band Structures of Simpler Metals . . . . .	267
VI. Excitonic Phases: Transcending Band Theory . . . . .	280
VII. Effect of Pressure on Transition Temperatures of Cooperative Phenomena . . . . .	290
VIII. Alloy Systems . . . . .	298
IX. Speculations on Crystal Structures in Extreme High Pressure Regime . . . . .	303
X. Concluding Remarks . . . . .	304
References . . . . .	305
Appendix I. Velocity of Sound in Metals and Metallic Alloys . . . . .	308
A. Non-linear plasma . . . . .	308
B. Extension of Bohr-Staver formula for liquid metal alloys . . . . .	309
Appendix II. Analogy between Excitonic Phase and Superconducting State . . . . .	311

CHAPTER 5	
Adsorption of Gases at High Pressures	
P. G. MENON	
I. Introduction . . . . .	313
II. Experimental Methods . . . . .	316
III. Results . . . . .	335
IV. Adsorption Equilibria in Multicomponent Gas Mixtures . . . . .	350
V. Miscellaneous . . . . .	359
References . . . . .	363
AUTHOR INDEX . . . . .	367
SUBJECT INDEX . . . . .	379

## CHAPTER 1

## High Pressure Mössbauer Studies†

H. G. DRICKAMER, S. C. FUNG and G. K. LEWIS, JR.‡

Department of Chemistry and Chemical Engineering, and Materials Research Laboratory, University of Illinois, Urbana, Illinois, U.S.A.

I. Introduction . . . . .	1
II. Pressure Effects . . . . .	11
A. Isomer shift . . . . .	11
B. Quadrupole splitting . . . . .	19
C. Magnetism . . . . .	23
D. The measurement of $f$ number . . . . .	26
E. Conversion of $\text{Fe}^{\text{III}}$ to $\text{Fe}^{\text{II}}$ . . . . .	29
F. High pressure Mössbauer studies on glass . . . . .	34
References . . . . .	38

## I. INTRODUCTION

The existence of the Mössbauer Effect (that is recoilless resonant radiation) was first discovered in 1958 (Mössbauer, 1958a, b). The principles involved have been reviewed in detail in the literature (Frauenfelder, 1963; Wertheim, 1964), so they will only be outlined here.

When a radioactive atom decays by gamma ray emission, the nucleus goes from an excited state to the ground state, and the energy of the gamma ray is a measure of the energy difference between these states. Consider first the case of a free atom. The emitted gamma ray will have associated with it a certain momentum, and in order to conserve momentum the atom must recoil. However, this recoil has associated with it kinetic energy which must reduce the energy of the gamma ray; the act of absorption of a gamma ray by a nucleus involves the inverse process. Thus the emission and absorption processes are not in resonance.

If, however, one fixes the atom in a crystal, the situation may be

† This work was supported in part by the United States Atomic Energy Commission under Contract AT(11-1)-1198. ‡ Present Address: Eastern Laboratories, E. I. duPont Company, Gibbstown, New Jersey, U.S.A.