

$(\partial S/\partial V)_T$  can be obtained from the experimental results by numerical differentiation. By using Mills & Grilly's  $p$ - $V$ - $T$  data and equation (7) we can then obtain the pressure at constant molar volume as a function of temperature, i.e. the isochores

$$p(T) = p_m - \int_T^{T_m} (\partial S/\partial V)_T dT, \quad (8)$$

$$p_m = p(T_m), \quad V = \text{const.}$$

The isochores are given in tables 6 and 7 for rounded values of the molar volume. The columns of these tables give immediately the isotherms, i.e.  $p = p(V)$  at constant temperature.




FIGURE 10. The lattice entropy of  $^3\text{He}$ . The numbered lines are lines of constant volume.

### 3.6.2. Compressibility

We have calculated the compressibility of solid  $^4\text{He}$  and  $^3\text{He}$  at  $0^\circ\text{K}$  from the  $0^\circ\text{K}$  isotherm

$$\beta = -\frac{1}{V} \left( \frac{\partial V}{\partial p} \right)_{T=0}. \quad (9)$$

$\beta$  is given at rounded values of molar volume in table 8.

### 3.6.3. Thermal expansion coefficient

The volume thermal expansion coefficient,  $\alpha$ , can be obtained from the thermodynamical relation

$$\alpha = \beta(\partial p/\partial T)_V. \quad (10)$$

$\alpha$  for solid  $^4\text{He}$  and  $^3\text{He}$  is given as a function of temperature and molar volume in table 9.