

$L_0$  is the latent heat of vaporization at  $0^\circ\text{K}$ ,  $p_m\Delta V_m$  is the work done in solidifying the liquid at  $0^\circ\text{K}$ , and the two integrals give the work of compression in the liquid and the solid range, respectively. ( $V_1$  is the volume of the liquid in equilibrium with its vapour,  $V_2$  in equilibrium with the solid, and  $V_3$  that of the solid in equilibrium with the liquid.)

TABLE 10

	$p$ (Kg/cm <sup>2</sup> )	$V$ (cm <sup>3</sup> /mole)	$U_0$ (cal/mole)
liquid $0^\circ\text{K}$	0	36.63	-5.04
liquid $0^\circ\text{K}$	29	26.0	-2.6
b.c.c. solid $0^\circ\text{K}$	29	24.2	-1.8




FIGURE 13. Energy relations at  $0^\circ\text{K}$  for solid  $^4\text{He}$  and  $^3\text{He}$ .  $U_z$  is the zero-point energy;  $U_0$  is the internal energy at absolute zero; and  $\Phi$  is the classical static lattice energy calculated from the de Boer-Michels potential. —, Experimental; ----, London (1954).

For  $^4\text{He}$  we have taken Swenson's (1950) estimate for one value of  $U_0$ . According to this,  $U_0$  for the solid at the melting pressure is  $-11.9$  cal/mole. In order to obtain  $U_0$  for smaller molar volumes we have to calculate the work of compression according to equation (11). This information is known from the present experiments between  $17.0$  and  $11.5$  cm<sup>3</sup>/mole. Up to  $V = 17$  cm<sup>3</sup>/mole we have used an extrapolation of the isotherm at  $0^\circ\text{K}$ . This extrapolation gives  $U_0 = -5.07$  cal/mole for  $V = 17.0$  cm<sup>3</sup>/mole.