

FIG. 2. The relative resistance of four rubidium samples as a function of temperature at low pressures (approximately 100 atm.). The resistances have been normalized to agree at 150° K. and then separated by intervals of 2 units to avoid overlapping.

MacDonald (1952) found an anomaly in the resistance of rubidium at about 180° K. and the present samples (except number 3) show this anomaly in various forms. In samples 1 and 2, the effect is quite evident, but in sample 4 the anomalous behavior takes the form of a gradual deviation from linearity with temperature which begins to be evident at about 150° K. Samples 1 and 2 showed marked thermal hysteresis in the neighborhood of the anomaly (cf. MacDonald 1952), the resistance measured with falling temperature being above that measured with rising temperature. To simplify the diagram only the falling temperature curves are shown here.

Sample number 3, which was heavily oxidized, shows little sign of the anomaly. (Its behavior is not shown above 260° K. since at this temperature premelting begins in this oxidized specimen and the resistance increases rapidly.) Other experiments in this laboratory (Hedgcock 1956) confirm that in highly oxidized rubidium the anomaly seems to be suppressed.

In addition to the work already referred to, Kelly and Pearson (1955) have investigated the anomaly in detail but at present its origin is still obscure.

The isothermal change in the neighborhood of room temperature with such a fine capillary pressure was appreciably larger than the resistivity change at 25° K. from Bridgman's measurements (Bridgman's measurements were on a single sample and have to be treated with caution because of compressibility.) This difference was observed in specimens melted into capillaries to demonstrate that the general qualitative picture of the anomaly at low temperatures.

In describing these results we shall proceed as follows:

- (1) measurements below 150° K.
- (2) measurements between 150° K. and 260° K.
- (3) measurement of the isothermal change in the neighborhood of room temperature.




FIG. 3. The resistivity of a rubidium sample as a function of temperature range.