


results in the form of relative
alkali metals at room tem-
peratures in this Figure is that in-
creases steadily with pressure and only
slightly with temperature.
By contrast the resistivity
decreases at low pressures and then rises
at all pressures in this range.



metals at 0° C. The curves show rela-

the behavior of these curves is as follows. No
marked change in temperature range under
conditions that the mean-square ampli-
tude increases monotonically with increasing
pressure in this pressure range. The lattice
effects cannot account for the minima
observed in the Li curve. These effects
change in K with volume. In
order to obtain relative values of K versus relative
volume in Fig. 24. (In order to obtain
relative values of K versus relative
volume has been estimated from
the data. It is clear that the main features
of these curves.




Fig. 24. K versus volume in the alkali metals deduced from the data in Fig. 23.

3. Comparison with Liquid Metals

The effect of pressure on the alkali metals in the liquid state has not been studied over such a wide range as for the solids. Bridgman has however made some measurements on the liquids. Of Cs he writes (Bridgman, 1949, p. 282): "Because of the location of the melting curve it was not possible to measure the resistance of the liquid metal at pressures high enough to reach the minimum [in the ρ - P curve], but simple extrapolation indicates that without much question the liquid will show the effect as well as the solid at temperatures above perhaps 140°, and there seems no reason to think that the mechanism responsible for the minimum has any essential connection with the lattice structure."

In Li, moreover, Bridgman finds that, as in the solid, the pressure coefficient of resistivity of the liquid is *positive* (in magnitude it is about 33% greater than that of the solid). In the other metals Bridgman finds negative pressure coefficients of resistivity of magnitude similar to those found in the corresponding solids.

To sum up what we know about the volume dependence of K : we know that: (1) at atmospheric pressure the sign of $\partial \ln K / \partial \ln V$ is different in different metals; (2) for the monovalent metals the sign of