

be tentative. With this reservation we would draw the following conclusions from our experiments. As far as the close-packed phases are concerned, solid helium in both isotopic forms appears to behave normally, i.e. it can be understood at least semi-quantitatively in terms of existing theories and concepts. For example, the zero-point energy can be estimated roughly by means of a crude cell model due to London, or by means of a quasi-harmonic model of the vibrations of the solids. Likewise the temperature dependence of the specific heats appears to be similar to that of a quasi-harmonic close-packed solid and the magnitude of the characteristic temperature at each density agrees with what is known about the mechanical properties of the solids.

The specific heat of the fluid isotopes can be understood qualitatively; at present there appears to be no possibility of a detailed comparison with fundamental theory.

The authors gratefully acknowledge the encouragement and interest of the late Dr D. K. C. MacDonald, F.R.S. They are grateful to Dr T. H. K. Barron for many valuable discussions; to Dr D. L. Martin for general advice on calorimetry; to Dr Barron and Dr A. M. Guénault for reading the manuscript; and to Dr C. A. Swenson for communicating his results before publication. The authors also wish to thank Mr A. A. M. Croxon and Mr R. L. Snowdon for their valuable technical help; and Miss B. A. Cotton and Mr D. R. Taylor for their work on the calculations.

REFERENCES

- Barron, T. H. K. & Klein, M. L. 1962 *Phys. Rev.* **127**, 1997.
 Bartholomé, E. & Eucken, A. 1937 *Trans. Faraday Soc.* **33**, 45.
 Beaumont, R. H., Chihara, H. & Morrison, J. A. 1961 *Proc. Phys. Soc.* **78**, 1462.
 Bernardes, N. 1960 *Phys. Rev.* **120**, 1927.
 Bernardes, N. & Brewer, D. F. 1962 *Rev. Mod. Phys.* **34**, 190.
 Clement, J. R. 1955 *Temperature, its measurement and control in science and industry*, vol. 2, p. 380. New York: Reinhold.
 de Boer, J. & Michels, A. 1938 *Physica*, **5**, 945.
 Domb, C. & Dugdale, J. S. 1957 *Progress in Low Temperature Physics*, vol. 2, p. 338. North Holland Publishing Company.
 Domb, C. & Salter, L. 1952 *Phil. Mag.* **43**, 1083.
 Dugdale, J. S. & Simon, F. E. 1953 *Proc. Roy. Soc. A*, **218**, 291.
 Eucken, A. 1916 *Verh. dtsch. Phys. Ges.* **18**, 4.
 Eucken, A. & Seekamp, H. 1928 *Z. phys. Chem.* **134**, 178.
 Franck, J. P. 1961 *Phys. Rev. Lett.* **7**, 435.
 Franck, J. P. & Martin, D. L. 1961 *Canad. J. Phys.* **39**, 1320.
 Grilly, E. R. & Mills, R. L. 1959 *Ann. Phys.* **8**, 1.
 Heltemes, E. C. & Swenson, C. A. 1961 *Phys. Rev. Lett.* **7**, 363.
 Heltemes, E. C. & Swenson, C. A. 1962 *Phys. Rev.* **128**, 1512.
 Hill, R. W. & Lounasmaa, O. V. 1960 *Phil. Trans. A*, **252**, 357.
 Hooton, D. J. 1955 *Phil. Mag.* **46**, 422, 433, 485.
 Hurst, R. P. & Levelt, J. M. H. 1961 *J. Chem. Phys.* **34**, 54.
 Keesom, W. H. 1942 *Helium*, p. 42. Amsterdam: Elsevier.
 Keesom, W. H. & Keesom, A. P. 1936 *Comm. Leiden*, no. 240b.
 Leighton, R. B. 1948 *Rev. Mod. Phys.* **20**, 165.
 Levelt, J. M. H. & Hurst, R. P. 1960 *J. Chem. Phys.* **32**, 96.
 London, F. 1954 *Superfluids*, vol. 2. New York: Wiley.