



REFERENCES

- [1] R. S. Krishnan. Proc. Indian Acad. Sci. A26 (1947) 432.
- [2] P. W. Bridgman, Phys. Rev. 38 (1931) 182; Proc. Am. Acad. Arts Sci. 76 (1945) 1.
- [3] G. K. Lewis, E. A. Perez-Albuerné and H. G. Drickamer, J. Chem. Phys. 45 (1966) 598.
- [4] C. W. Garland and R. Renard, Critical phenomena (U.S. Natl. Bur. Stds., Misc. Publ. No. 273, 1966) p. 202.
- [5] E. L. Wagner and D. F. Horing, J. Chem. Phys. 18 (1950) 296 and references cited therein.
- [6] J. F. Asell and M. Nicol, J. Chem. Phys. 49 (1968) 5395.
- [7] Y. Ebisuzaki, W. D. Ellenson and M. Nicol, in preparation.
- [8] C. Postmus, J. R. Ferraro and S. S. Mitra, Phys. Rev. 174 (1968) 983.
- [9] H. S. Gutowsky, G. E. Pake and R. Bersohn, J. Chem. Phys. 22 (1954) 643.
- [10] P. S. Leung, T. I. Taylor and W. W. Havens Jr., J. Chem. Phys. 48 (1968) 4912.
- [11] T. Naganiya, Compt. Rend. Reunion Ann. Comm. Thermodynam., Union Intern. Physics (Paris) (1952) 251.
- [12] K. Nakamoto, M. Margoshes and R. E. Rundle, J. Am. Chem. Soc. 77 (1955) 6840.
- [13] J. P. Mathieu and H. Poulet, Spectrochim. Acta 16 (1960) 696.
- [14] Y. Ebisuzaki and M. Nicol, to be published.



the slopes of the
-1/kbar⁻¹.



and transverse
are. The slopes
upper); -0.29
r⁻¹.

ts along the
the frequen-
can be attri-
hydrogen bond

he ν_4 mode is
longitudinal
transverse com-
pressure range
clear at pres-
nancing site
l intensity ef-
will be de-
ether with a
pressure on the
nds of the spec-
aminary results
ous assignment