

the results  
2, are next

ement eigen-  
odes,  $A^9$  and  
d  $-5.2^\circ$  for  
on directions  
is favorable  
ing rise to the  
e oscilloscope.  
the 3-mm-  
rs. The cor-  
e  $+14.6^\circ$  for  
se deviations  
our egregious  
as in the Y-Z  
for bismuth.

flux, or ray  
or our dimen-  
e reaching the  
the energy is  
to the pulses  
responds with  
in relation to  
y not enough  
ction to reach  
to the trans-

ion angle, the  
e echoes, and  
every mode.  
axis, predicted  
rock salt and  
uth, did not  
cho patterns.  
n  $28^\circ 40'$ ; for

her the 3-mm-  
nder-receiver.  
arger one con-  
displays ob-  
arently upon

the effects of  
nce we do not  
nen geometry.  
ned effects of  
on are not as  
LR's velocities  
his antimony  
e for bismuth.

sed on plane  
specimens are

).




FIG. 2. Energy density flow, displacement, propagation and pure-mode directions in Y-Z mirror plane.  $\delta_K^s$  is the angle between the vectors  $K$  and  $s$  where  $s$  is any of the vectors  $A^9$  and  $P^9$ . The signs for  $\delta_K^s$  indicate placement at opposite sides of  $K$  with  $+\delta$  counterclockwise. Parts (a) and (b) are for antimony, (c) and (d) for bismuth; (a) and (c) are for the  $45^\circ$  modes, and (b) and (d) for the  $135^\circ$  modes.

of finite dimensions. For isotropic circular bars the dilatational and distortional wave phase velocities have been shown by Pochhammer<sup>20,21</sup> to depend on the ratio of the cross section radius to wavelength  $a/\lambda$  and upon two functions of the Lamé stiffness constants. Chree<sup>22</sup> extended Pochhammer's results to noncircular, normal

cross-sectioned cylinders and to nonisotropic media. In the absence of an exact treatment giving the longitudinal and two transverse phase velocities in anisotropic cubes it is reasonable to assume that the size of the correction for each phase velocity would be different. If these corrections are large in relation to the experimental errors, fitting the plane-wave formalism of redundancy eight to the 14 corrected velocities is not assured. That we are able to do so, however, suggests that the corrections are not significant. Our large minimum value of about 25 for  $k/\lambda$ , where  $k$  is the radius

<sup>20</sup> A. E. H. Love, *A Treatise on the Mathematical Theory of Elasticity* (Dover Publications, Inc., New York, 1944), p. 287.

<sup>21</sup> H. Kolsky, *Stress Waves in Solids* (Dover Publications, Inc., New York, 1963), p. 54.

<sup>22</sup> A. E. H. Love, *A Treatise on the Mathematical Theory of Elasticity* (Dover Publications, Inc., New York, 1944), p. 290.