

## REFERENCES

- [1] H.A.Brouwer and C.G.Egeler, The glaucophane schist facies metamorphism in the schistes lustres nappe of Corsica, Proc. Kon. Ned. Akad. Wetensch. tw. rks. 58 (1952) 3.
- [2] M.A.Autran, Description de l'association jadeite + quartz et des paragénèses minérales associées, dans les schistes lustres de Sant Andrea di Cotone (Corse), Bull. Soc. Franc. Min. Crist. 87 (1964) XLIII-XLIV.
- [3] H.G.F.Winkler, Petrogenesis of Metamorphic Rocks (Springer-Verlag, New York, 1965) pp. 142, 151.
- [4] R.G.Coleman, Composition of jadeitic pyroxene from the California metagreywackes, U.S. Geol. Surv. Prof. Paper 525.C (1965) C25-C34.
- [5] R.G.Coleman and Joan R.Clark, Pyroxenes in the blueschist facies of California, Am. J. Sci. 266 (1968) 43.
- [6] E.J.Essene and W.S.Fyfe, Omphacite in Californian metamorphic rocks, Contr. Mineral. Petrol. 15 (1967) 1.
- [7] R.C.Newton and J.V.Smith, Investigations concerning the breakdown of albite at depth in the earth, J. Geol. 75 (1967) 268.
- [8] I.Kushiro, Clinopyroxene solid solutions at high pressures, in: Ann. Rept. Director, Geophys. Lab. for 1964-1965, Carnegie Inst. Wash. Yearbook 64 (1965) 112.
- [9] F.J.Turner and J.Verhoogen, Igneous and Metamorphic Petrology, 2nd ed. (McGraw-Hill, New York, 1960) 544.
- [10] E.J.Essene, An occurrence of cymrite in the Franciscan formation, California, Am. Mineral. 52 (1967) 1885.