

TABLE 4

Chemical composition of glass bombs (*Fladen*) from suevite, collected at 9 different localities in the Ries basin (Amerbach, Aufhausen, Aumühle, Bollstadt, Doosweiher, Fünfsetten, Heerhof, Otting, Schmähingen).

	9 unrecrystallized glass bombs		17 recrystallized glass bombs	
	Average composition	Standard deviation	Average composition	Standard deviation
SiO ₂	63.54	1.03	64.04	1.15
TiO ₂	0.81	0.08	0.78	0.10
Al ₂ O ₃	15.10	0.43	15.28	0.80
Fe ₂ O ₃	0.99	0.19	1.42	0.72
FeO	3.75	0.19	2.39	1.26
MnO	0.10	0.01	0.08	0.03
MgO	2.71	0.19	1.71	0.92
CaO	3.45	0.34	3.98	0.54
Na ₂ O	2.86	0.29	3.59	0.68
K ₂ O	3.71	0.16	3.50	0.76
P ₂ O ₅	0.36	0.12	0.32	0.08
H ₂ O ⁺	2.73	0.30	2.72	1.05
CO ₂	0.37	0.22	0.33	0.19
Total	100.48	—	100.14	—
Fe ²⁺ /total Fe	0.81	—	0.65	—

consists of fine schlieren which differ in refractive index and in color. The structure reflects the heterogeneous nature of the parent rock, the intense movement of the melt, and the short duration of the molten state (Fig. 7).

The chemical composition of the *Fladen* is very uniform. Table 4 shows the average chemical composition of 9 unrecrystallized and 17 re-




Fig. 7. Unrecrystallized molten glass, from the suevite at Otting. Schlieren, circular vesicles, angular mineral fragments, and fused silica (lechatelierite) (light-colored elongate areas in upper left) are all present.

crystallized *Fladen*, collected at 9 different localities around the Ries basin. As can be seen from the standard deviations, the scattering is relatively low. The composition of the *Fladen* is consistent with the assumption that they were produced by melting of the granitic basement rocks. Their chemical composition differs appreciably from the composition of *Moldavite* tektites, and theories which derive the Moldavites from melt produced by the Ries impact event have to explain this discrepancy (Engelhardt and Hörz, 1965). The *Fladen* differ in their cooling histories. There are rapidly chilled *Fladen* which contain clear, undevitrified glass; specimens of more slowly cooled *Fladen* show different types of devitrification. Cristobalite, feldspar, and pyroxene are the main devitrification products; they crystallized after movement in the melt had ceased. This conclusion follows from the observations that orientation of the new crystals is not affected by the flow pattern of the glass and that there are no parallel arrangements of elongate crystals of the type so familiar in lava flows. Furthermore, it is certain that devitrification took place after deposition of the suevite breccia, because it can be seen that, in

some suevite quarries, the lowest, rapidly cooled layer of suevite contains only unrecrystallized glasses, whereas higher in the suevite, all the *Fladen* are devitrified. In the quarry of Otting, for instance, the chilled basal layer is about 1 m thick. Chilled layers with unrecrystallized glasses at the bottom and the top of suevite have also been found in the bore hole at Wörnitzostheim (Förstner, 1967).

ACKNOWLEDGMENT

The authors thank the Deutsche Forschungsgemeinschaft for financial support.

REFERENCES

- Ahrens, T. J. and J. T. Rosenberg, Shock metamorphism: experiments on quartz and plagioclase, *this vol.*, p. 59.
- Baranyi, I., Sedimentary rock fragments in the suevite breccia, Diplom. thesis, Tübingen, 1966.
- Bunch, T. E., A. J. Cohen, and M. R. Dence, Shock-induced structural disorder in plagioclase and quartz, *this vol.*, p. 509.
- Burri, C., R. L. Parker und W. Wenk, Die optische Orientierung der Plagioklase. Unterlagen und Diagramme zur Plagioklasbestimmung nach der Drehtischmethode, Basel, 1967.
- Chao, E. C. T., Shock effects in certain rock-forming minerals, *Science*, 156, 192-202, 1967.
- Chao, E. T. C. and J. Littler, Additional evidence for the impact origin of the Ries basin, Bavaria, Germany, *Geol. Soc. Am. Abstr. for 1962*, 127, 1963.
- Cummings, D., Kink-bands: shock deformation of biotite resulting from a nuclear explosion, *Science*, 148, 950-952, 1965.
- David, E., Flight of tektites from meteorite impact, *Z. f. Naturforsch.* 21a, 1133-1137, 1966.
- Engelhardt, W. v., Chemical composition of Ries glass bombs, *Geochim. Cosmochim. Acta*, 31, 1677-1689, 1967.
- Engelhardt, W. v. and F. Hörz, Riesgläser und Moldavite, *Geochim. Cosmochim. Acta* 29, 609-620, 1965.
- Engelhardt, W. v., J. Arndt, D. Stöffler, W. F. Müller, H. Jeziorkowski and R. A. Gubser, Diaplektische Gläser in den Breccien des Ries von Nördlingen als Anzeichen für Stosswellenmetamorphose, *Contr. Mineral. and Petrol.*, 15, 93-102, 1967.
- Engelhardt, W. v., F. Hörz, D. Stöffler and W. Bertsch, Observations on quartz deformation in breccias of West Clearwater Lake, Canada and the Ries basin, Germany, *this vol.*, p. 475.
- Förstner, U., Petrographische Untersuchungen des Suevit aus den Bohrungen Deiningen und Wörnitzostheim im Ries von Nördlingen, *Contr. Mineral. and Petrol.*, 15, 281-308, 1967.
- Griggs, D. T., F. J. Turner and H. C. Heard, Rock deformation, *Geol. Soc. Am. Mem.* 79, 39-104, 1960.
- Hörz, F., Untersuchungen an Riesgläsern, *Beitr. Mineral. und Petrogr.* 11, 621-661, 1965.
- Milton, D. J. and P. S. De Carli, Maskelynite: formation by explosive shock, *Science*, 140, 670-671, 1963.
- Müller, W. F., Unpublished investigations on the action of shock waves on quartz and feldspar, Mineral. Inst. Univ. Tübingen, Publication in preparation, 1967.
- Rinehart, J. S., Intense destructive stresses resulting from stress wave interactions, *this vol.*, p. 433.
- Robertson, P. B., M. A. Vos and M. R. Dence, Deformation in rock forming minerals from Canadian craters, *this vol.*, p. 31.
- Seeger, A., Kristallplastizität, in Flüge, F., *Handbuch der Physik*, VII/2, Kristallphysik II, Berlin, Göttingen, Heidelberg, p. 1-210, 1958.
- Shoemaker, E. M. and E. C. T. Chao, New evidence for the impact origin of the Ries basin, Bavaria, Germany, *J. Geophys. Res.* 66, 3371-3378, 1961.
- Short, N. M., Effects of shock pressures from a nuclear explosion on mechanical and optical properties of granodiorite, *J. Geophys. Res.* 71, 1195-1217, 1966.
- Stöffler, D., Anzeichen besonderer mechanischer Beanspruchung an Mineralien der Kristallineinschlüsse des Suevits (Stosswellenmetamorphose), *N. Jb. Miner. Mh.*, 9-11, 350-354, 1965.
- Stöffler, D., Zones of impact metamorphism in the crystalline rocks of the Nördlinger Ries crater, *Contr. Mineral. and Petrol.* 12, 15-24, 1966.
- Stöffler, D., Deformation und Umwandlung von Plagioklas durch Stosswellen in den Gesteinen des Nördlinger Ries, *Contr. Mineral. and Petrol.*, 16, 51-83, 1967.
- Wackerle, J., Shock compression of quartz, *J. Appl. Phys.* 33, 922-937, 1962.