

REFERENCES

1. D. B. McWhan and M. P. Sarachik, Bull. Am. Phys. Soc. 12, 504 (1967).
2. E. Fawcett and G. K. White, J. Appl. Phys. 39, 576 (1968).
The measurements reported in this reference were on a sample of Pd which was later found to contain several hundred ppm Fe impurity, and which exhibited a strongly temperature and field-dependent susceptibility. The present sample of Pd was made from Matthey-Bishop wire containing 4 ppm Fe and < 10 ppm total metallic impurities, and its susceptibility was field-independent and essentially independent of temperature at low temperatures. We are indebted to E. Bucher for these susceptibility measurements.
3. N. D. Lang and H. Ehrenreich, J. Appl. Phys. 38, 1316 (1967); Phys. Rev. (to be published).
4. S. Doniach and E. P. Wohlfarth, Proc. Roy. Soc. A296, 442 (1967).
5. D. J. Kim, Phys. Rev. 149, 434 (1966).
6. Similar non-linear magnetostriction is observed even at temperatures well below the Curie temperature in an alloy $\text{Pt}_{99}\text{Fe}_{01}$ where the magnetostriction varies as the square of the magnetization (E. Fawcett and R. C. Sherwood, to be published), in accordance with the prediction of a Stoner model for a weak itinerant ferromagnet (P. Wohlfarth, to be published).