

KRAUT-KENNEDY MELTING RELATIONSHIP AT HIGH PRESSURES EXTENDED TO CERTAIN PHASE TRANSFORMATIONS *

R. B. FISCHER

Rocky Flats Division, Dow Chemical Company, Golden, Colorado, USA

Received 8 November 1966

Kraut and Kennedy¹⁾ have presented data which suggest that if a solid is compressed isothermally at room temperature and then heated at constant pressure until it melts, the additional energy required to melt the solid at higher pressure depends *linearly* on the amount by which the solid was compressed. A natural extension of this law for melting points would seem to apply to some phase transformations involving volume increases.

In a preliminary examination, experimental data for the alpha to beta transformation in plutonium^{2, 3)} and for the alpha to beta transformation in uranium^{4, 5)} were plotted with the results as shown in fig. 1. Reasonable linearity is noted in both cases.

Thus one might use the transformation temperature formulas:

$$T_{\text{Pu } \alpha \rightarrow \beta} (\text{°C}) = 120 (1 + 49.3 \Delta V/V_0), \quad (1)$$

and

$$T_{\text{U } \alpha \rightarrow \beta} (\text{°C}) = 660 (1 + 7.8 \Delta V/V_0), \quad (2)$$

where $\Delta V/V_0$ is the isothermal compression at room temperature.

The limit of (1) is uncertain at present, and (2) holds only to 40 kbar where a triple point⁵⁾ has been found.

More experimental pressure-temperature data are needed on other systems for further verification of the Kraut-Kennedy relationship.

* Work performed under U.S. Atomic Energy Commission Contract AT(29-1)-1106.




Fig. 1

References

- 1) E. A. Kraut and G. C. Kennedy, Phys. Rev. Lett. **16** (1966) 608
- 2) P. W. Bridgman, J. Appl. Phys. **30** (1959) 214
- 3) D. R. Stephens, J. Phys. Chem. Solids **24** (1963) 1197
- 4) P. W. Bridgman, Proc. Am. Acad. Arts Sci. **76** (1948) 75
- 5) W. Klement, Jr., A. Jayaraman and G. C. Kennedy, Phys. Rev. **129** (1963) 1971