

lesser concentration dependent than for the other methods. The Bjerrum distance appears to give K values slightly different from the Davies-Otter-Prieur equation used with lattice⁶. The Davies-Otter-Prieur results of Spedding and Jackson⁶, in agreement with the results of the other solution, in the neighborhood of 2.4×10^{-4} at low concentration, is seen from Table V that regardless which method is used, K_m^+ at atmospheric pressure approaches values in the original data for $\text{La}^{2+}(\text{SO}_4)^-$ solution are shown in Table VIII.

$\text{La}^{2+}\text{SO}_4^+$ of 23.2 is used rather than the value of 40.0 proposed by Spedding and Jackson because this value yielded values of K_m^+ which showed less concentration dependence of SO_4^{2-} concentration.

Discussion

It is seen from Table V that regardless which method dependence. The pressure dependence of the $\text{La}^{2+}\text{SO}_4^+$ dependence for pressure dependence as for A_b^+ for $\text{La}^{2+}(\text{SO}_4)^-$. ion was taken to be the same as for A_b^+ . Results for pressure dependence of equilibrium concentration are shown in Table VII. Results for pressure dependence of equilibrium concentration are shown in Table VII. The original data distribution of equivalent con-

P, atm		$\text{La}^{2+}(\text{SO}_4)^-$		La^{2+}		SO_4^{2-}		$\text{La}^{2+}\text{SO}_4^+$	
atm	0.2	0.3	0.3	0.6	1.2	2.4	3.1	8.2	mequiv/l.
atm	0.2	0.3	0.3	0.6	1.2	2.4	3.1	8.2	mequiv/l.
atm	0.2	0.3	0.3	0.6	1.2	2.4	3.1	8.2	mequiv/l.
atm	0.006	0.012	0.040	0.12	0.32	0.43	1.5	0.006	^a See footnote a of Table IV.
atm	0.006	0.013	0.041	0.12	0.33	0.45	1.6	0.006	^b See footnote b of Table IV.
atm	0.007	0.014	0.045	0.13	0.36	0.49	1.8	0.007	for aqueous $\text{La}^{2+}(\text{SO}_4)^-$ at 25°.
atm	0.007	0.015	0.049	0.14	0.37	0.50	1.7	0.007	Table V: Dissociation Constant $K_m (\times 10^4)$
atm	0.007	0.016	0.052	0.15	0.40	0.54	1.9	0.007	of $\text{La}^{2+}\text{SO}_4^+$.
atm	0.008	0.018	0.057	0.15	0.40	0.54	1.8	0.008	See a function of Pressure as a function of Pressure.
atm	0.008	0.019	0.058	0.16	0.41	0.56	1.9	0.008	Table VI: - ΔV° (ml/mole) for aqueous $\text{La}^{2+}(\text{SO}_4)^-$ at 25°.
atm	0.009	0.020	0.061	0.17	0.43	0.59	2.0	0.009	Table IV: Concentration ($\times 10^3$, equiv/l.) of $\text{La}^{2+}(\text{SO}_4)^-$ at 25°.
atm	0.011	0.023	0.069	0.18	0.46	0.61	2.0	0.011	of $\text{La}^{2+}\text{SO}_4^+$ in aqueous $\text{La}^{2+}(\text{SO}_4)^-$ at 25°.
atm	0.011	0.024	0.070	0.18	0.47	0.63	2.1	0.011	^c Data obtained using the basic
atm	0.011	0.025	0.073	0.19	0.48	0.65	2.1	0.011	equation with $d = 3.6$ A for the 2-1 case.
atm	0.015	0.030	0.085	0.22	0.54	0.72	2.3	0.015	^d Data obtained using the Davies-Otter-Pri
atm	0.015	0.031	0.088	0.22	0.54	0.72	2.3	0.015	conductance equation, $d = 5$ A for the 3-2 case and
atm	0.015	0.030	0.086	0.22	0.53	0.70	2.2	0.015	$Z\Delta Z^2/2eKT$, calculated from Birrums equation, $d = 3.6$ A for the 2-1 case.
atm	0.015	0.030	0.085	0.22	0.52	0.69	2.2	0.015	Data obtained using the Davies-Otter-Pri

Table VIII: Copy of Original Conductivity Data for Aqueous Solutions of $\text{La}_2(\text{SO}_4)_3$ at 25°; Teflon Cell with Pyrex Bar between Electrodes: Cell Constant 0.457, Measured at 0.02 M KCl

P , atm	10 ³ equiv/l. of $\text{La}_2(\text{SO}_4)_3$							H ₂ O
	0.2046	0.3072	0.6144	1.206	2.412	3.072	8.190	
Conductivity in μmhos								
1	53.88	76.31	131.2	217.1	361.2	433.4	880.7	2.31
250	56.73	80.81	140.3	233.9	391.6	470.8	960.0	2.70
500	59.16	84.69	148.4	248.9	419.5	505.1	1033.5	3.11
750	61.22	88.08	155.3	262.2	444.6	536.7	1103.8	3.59
1000	63.01	90.87	161.1	274.1	467.6	565.3	1171.0	4.09
1250	64.53	93.35	166.4	284.5	488.1	590.9	1229.8	4.64
1500	65.85	95.45	170.7	293.2	505.7	613.5	1284.5	5.26
1750	66.94	97.18	174.4	300.8	521.2	633.5	1332.3	5.88
2000	67.85	98.73	177.6	307.2	534.7	650.9	1376.8	6.52
1 ^a	54.11	76.42	131.4	217.1	361.3	433.8	883.8	2.59

^a Readings taken the day after the pressure run.

At atmospheric pressure there is at most only a 5% difference in the ΔV° values obtained by the three methods. At the highest pressures and highest concentration the largest difference in the ΔV° values occur.

The atmospheric pressure values of ΔV° are of the same order as observed for NH_4OH ¹¹ and organic solutions¹² and very close to the value of -23.4 ml/mole calculated by Owen and Brinkley^{2a} for water. It is not known if a possible multistate configuration^{13,14} exists similar to that of MgSO_4 or MnSO_4 or if there exists only one form of the $(\text{LaSO}_4)^+$ ion pair. There is some indication that the rare earth sulfates show large ultrasonic absorption¹⁵ but until detailed experimental results are available, it is not possible to make an interpretation incorporating acoustic data.

Based on values of partial molal volume assigned by Owen and Brinkley^{2a} to La^{3+} of -38.3 ml/mole and SO_4^{2-} of +14.5, the partial molal volume of the $(\text{LaSO}_4)^+$ ion pair at atmospheric pressure and at the lowest concentration varies from -0.9 to +0.2 ml/mole depending upon the method used to calculate theoretical values of equivalent conductance.

(11) S. D. Hamann, "Physico-Chemical Effects of Pressure," Academic Press, New York, N. Y., 1957, p 151.

(12) A. Distechi and S. Distechi, *J. Electrochem. Soc.*, **112**, 350 (1965).

(13) F. H. Fisher, *J. Phys. Chem.*, **69**, 695 (1965).

(14) G. Atkinson and S. K. Kor, *ibid.*, **69**, 128 (1965).

(15) G. Atkinson, private communication.