

[Reprinted from the Journal of Physical Chemistry, 71, 819 (1967).]
 Copyright 1967 by the American Chemical Society and reprinted by permission of the copyright owner.

Effect of Pressure on the Dissociation of the $(\text{LaSO}_4)^+$ Complex Ion¹

by Frederick H. Fisher and Douglas F. Davis

University of California, San Diego, Marine Physical Laboratory of the Scripps Institution of Oceanography,
 San Diego, California 92152 (Received June 30, 1966)

Electrical conductance as a function of pressure up to 2000 atm has been measured on aqueous solutions of lanthanum sulfate at 25° at seven concentrations from 0.0002 to 0.0082 equiv/l. The effect of pressure on the dissociation constant of the $(\text{LaSO}_4)^+$ complex ion pair was calculated using three different methods. At atmospheric pressure ΔV° ranged from -21.2 to -26.2 ml/mole and at 2000 atm ΔV° ranged from -6.8 to -11.8 ml/mole depending on the method used and concentration. Based on the atmospheric pressure value of $\bar{V}_2^\circ = -23.8$ for $\text{La}_2(\text{SO}_4)_3$ the partial molal volume of the $(\text{LaSO}_4)^+$ ion at the lowest concentration is -0.9 to 0.2 ml/mole.

The very small partial molal volume assigned by Owen and Brinkley^{2a} to the La^{3+} ion, $\bar{V}_i^\circ = -38.3$ ml/mole relative to the H^+ ion for which $\bar{V}_i^\circ = 0$ ml/mole, suggested to F. H. F. that there might be an even greater effect of pressure on $\text{La}_2(\text{SO}_4)_3$ than was observed for MgSO_4 ^{2b} or MnSO_4 .³

Measurements of the electrical conductivity were made on aqueous solutions of $\text{La}_2(\text{SO}_4)_3$ at 25° in essentially the same manner as for MgSO_4 . It was possible to use Pyrex connected electrodes throughout the measurements with a small correction (10^{-6} atm⁻¹) for the coefficient of linear compression of Pyrex.⁴

Lanthanum chloride was prepared from Lindsay lanthanum oxide in the manner of Nathan,⁵ while the lanthanum sulfate purchased from K and K Laboratories was recrystallized once, also after Nathan. Reagent grade potassium chloride and potassium sulfate were obtained from Mallinckrodt and Matheson Coleman and Bell, respectively. All solutions were within ± 0.2 pH unit of the conductance water (pH 5.8).

Results

Ratios of equivalent conductivities as a function of pressure are shown in Table I. By making use of earlier data^{6,7} obtained at 1 atm, values for equivalent conductance as a function of pressure were calculated using the ratios in Table I and are shown in Table II. In Table III measurements to 2000 atm on KCl solutions obtained in the 30-ml cylindrical Teflon cell show agreement to within 0.1% with those obtained in a 12-cc spherical Pyrex cell.

Table I: Λ_p/Λ_1 for Aqueous Solutions of $\text{La}_2(\text{SO}_4)_3$ at 25°

P, atm	mequiv/l.						
	0.2	0.3	0.6	1.2	2.4	3.1	8.2
250	1.036	1.044	1.056	1.064	1.072	1.074	1.078
500	1.063	1.078	1.102	1.119	1.135	1.139	1.147
750	1.082	1.106	1.140	1.166	1.190	1.198	1.213
1000	1.096	1.125	1.169	1.206	1.239	1.249	1.274
1250	1.104	1.140	1.193	1.239	1.292	1.296	1.326
1500	1.107	1.149	1.209	1.263	1.314	1.330	1.372
1750	1.107	1.153	1.222	1.283	1.342	1.361	1.411
2000	1.103	1.156	1.231	1.299	1.365	1.387	1.447

If c is concentration of the solute in equivalents, then the dissociation constant for the reaction



may be expressed as follows⁸

$$K = [(c/3 - x)(c/2 - x)/x]\pi^f \quad (2)$$

(1) This paper represents results of research sponsored by the Office of Naval Research.

(2) (a) B. B. Owen and S. R. Brinkley, Jr., *Chem. Rev.*, **29**, 461 (1941); (b) F. H. Fisher, *J. Phys. Chem.*, **66**, 1607 (1962).

(3) F. H. Fisher and D. F. Davis, *ibid.*, **69**, 2595 (1965).

(4) L. H. Adams, *J. Am. Chem. Soc.*, **53**, 3780 (1931).

(5) C. C. Nathan, W. E. Wallace, and A. L. Robinson, *ibid.*, **65**, 790 (1943).

(6) F. H. Spedding and S. Jaffe, *ibid.*, **76**, 882 (1954).

(7) I. L. Jenkins and C. B. Monk, *ibid.*, **72**, 2695 (1950).

(8) At atmospheric pressure the difference between molar and molal units will be neglected.

Table II: Λ_p , Equivalent Conductance for Aqueous $\text{La}_2(\text{SO}_4)_3$ Solutions at 25°

P , atm	mequiv/l.						
	0.2	0.3	0.6	1.2	2.4	3.1	8.2
1	119.0 ^c	109.5 ^c	94.7 ^c	80.7 ^a	67.1 ^b	63.3 ^b	48.8 ^b
250	123.3	114.3	100.0	85.9	71.9	68.0	52.6
500	126.5	118.0	104.4	90.3	76.2	72.1	56.0
750	128.8	121.1	108.0	94.1	79.8	75.8	59.2
1000	130.4	123.2	110.7	97.3	83.1	79.1	62.2
1250	131.4	124.8	113.0	100.0	86.7	82.0	64.7
1500	131.7	125.8	114.5	101.9	88.2	84.2	67.0
1750	131.7	126.3	115.7	103.5	90.0	86.2	68.9
2000	131.3	126.6	116.6	104.8	91.6	87.8	70.6

^a I. L. Jenkins and C. B. Monk, *J. Am. Chem. Soc.*, **72**, 2695 (1950). ^b F. H. Spedding and S. Jaffe, *ibid.*, **76**, 882 (1954).

^c Interpolated graphically from combined data of *a* and *b*.

Table III: Comparison of KCl Conductance Ratios Λ_p/Λ_1 for Two Cells at 25.00 ± 0.02° and 10-kc Bridge Frequency

P , atm	Teflon cell ^a	Glass cell ^b
250	1.0090	1.0094
500	1.0145	1.0147
750	1.0175	1.0178
1000	1.0177	1.0180
1250	1.0167	1.0168
1500	1.0133	1.0134
1750	1.0086	1.0087
2000	1.0028	1.0025

^a Λ_p/Λ_1 average over five concentrations of KCl from 0.0005 to 0.02 *M* measured in 30-ml cylindrical Teflon cell with parallel Pyrex-spaced platinum electrodes coated with platinum black.

^b Λ_p/Λ_1 for 0.02 *M* KCl measured in 12-ml spherical Pyrex cell with parallel shiny platinum electrodes.

where x is the concentration in equivalents of the $(\text{LaSO}_4)^+$ complex ion and π^f is the activity coefficient product.

Calculations were made using the mixture rule in which the solution is regarded as a mixture⁹ of a 1-2 salt, $(\text{LaSO}_4)_2\text{SO}_4$, at equivalent concentration x and the 3-2 salt, $\text{La}_2(\text{SO}_4)_3$, at equivalent concentration $c - 3x$. The observed equivalent conductance of the solution is written then as

$$\Lambda_{\text{obs}} = (x/c)\Lambda_{12} + (c - 3x)/c\Lambda_{23} \quad (3)$$

where the Λ_{12} and Λ_{23} are calculated from theory and x is solved for by successive approximations. The activity coefficient product is⁶

$$\pi^f = [(f_{32\pm}^{\pm})^{5/2}/(f_{12\pm}^{\pm})^{3/2}] \quad (4)$$

where

$$-\log f_{ij\pm} = (A|z_i z_j| \sqrt{I}) / (1 + B\tilde{a} \sqrt{I}) \quad (5)$$

and the ionic strength $I = 2.5c - 6x$.

Calculations of x and K were made using three different methods as follows.

Method a: The Davies-Otter-Prue equation with appropriate modifications for pressure-dependent terms^{2b}

$$\Lambda = \Lambda^\circ - [(R\Lambda^\circ / \{1 + B\tilde{a} \sqrt{I}/2\}) + E] \times (\sqrt{I} / \{1 + B\tilde{a} \sqrt{I}\}) \quad (6)$$

was used to calculate the equivalent conductance. Bjerrum distances (at atmospheric pressure) of 21.4 Å for the 3-2 and 7.14 Å for the 1-2 salt were used for \tilde{a} and the same pressure dependence was applied as in the MgSO_4 work.

Method b: Equation 6 was used but distances of $\tilde{a} = 5$ Å for the 3-2 and $\tilde{a} = 3.6$ Å for the 1-2 salt were used; these are the same values that Spedding and Jaffe used.

Method c: The Onsager equation⁹

$$\Lambda = \Lambda^\circ - [R\Lambda^\circ + E] \sqrt{I} \quad (7)$$

was used to calculate the equivalent conductance and the \tilde{a} distances of 5 and 3.6 Å were used in the activity coefficient calculation.

The pressure-dependent forms of eq 6 and 7 were used to calculate x and K as a function of pressure in a manner similar to that described earlier,^{2b} and the results for x in molar units are shown in Table IV. The molal dissociation constant, K_m , shown in Table V was obtained by dividing the values of K in molar units by the ratio of the density of water at pressure P to that at atmospheric pressure.

From the equation^{2a}

$$(\partial \ln K_m / \partial p)_{T,m} = -(\Delta V^\circ / RT) \quad (8)$$

values of ΔV° are calculated corresponding to the three different methods and are shown in Table VI.

The pressure dependence of infinite dilution equivalent conductivity, Λ_p° , for the $\text{La}_2(\text{SO}_4)_3$ was determined from the equation

$$\Lambda_p^\circ [\text{La}_2(\text{SO}_4)_3] = \Lambda_p^\circ [\text{LaCl}_3] + \Lambda_p^\circ [\text{K}_2\text{SO}_4] - \Lambda_p^\circ [\text{KCl}] \quad (9)$$

The Λ_p° values for the La^{3+} and SO_4^{2-} ions were calculated using the KCl transference number data of Wall and Gill.¹⁰ It was assumed that their transference number data to 1000 atm could be extrapolated linearly to 2000 atm. The Jenkins and Monk⁷ value for

(9) C. W. Davies, "Ion Association," Academic Press, New York, N. Y., 1962.

(10) F. T. Wall and S. J. Gill, *J. Phys. Chem.*, **59**, 278 (1955).