ressed by the

(3)

of atoms which ion which leads n volume of nately 2-3 is g the estimated

rrelation bethe present losely related future meaa larger numcomplete picals. measure the dered as well since the mag- 10^{-13} cm²/sec red would be diffusion ex-An alloy sysements in both

d be possible

as the degree

atio $\Delta V/V_M$

the activation -equiatomic :m3/mole or at 944.5°C and This value of orted value of dicates that quite similar mic metal. or D. Lazarus D. Gupta and wing access to lication.

Warschauer

1969). ity of Illinois,

ol. Phys. 41,

hysics, 3rd

AIME 175, 742

und S. Kachi,

11 D. N. Yoon and R. N. Jeffery, J. Phys. Chem. Solids (to be published).

12 N. H. Nachtrieb, J. A. Weil, E. Catalano, and A. W. Lawson, J. Chem. Phys. 20, 1189 (1952).

13R. A. Hultsch and R. G. Barnes, Phys. Rev. 125, 1832 (1962).

14R. E. Hanneman, R. E. Ogilvie, and H. C. Gatos, Trans. Met. Soc. AIME 233, 691 (1965).

15E. D. Albrecht and C. T. Tomizuka, J. Appl. Phys.

35, 3560 (1964). ¹⁶D. Gupta and R. N. Jeffery (unpublished).

¹⁷D. N. Yoon and D. Lazarus, Bull. Am. Phys. Soc. 15, 381 (1970); D. Lazarus, D. N. Yoon, and R. N. Jeffery, Europhysics Conference on Atomic Transport in Solids and Liquids, Marstrand, Sweden, 1970 (unpublished).

18M. Beyeler and Y. Adda, J. Phys. (Paris) 29, 345 (1968); B. M. Butcher, H. Hutto, and A. L. Ruoff, Appl. Phys. Letters 7, 34 (1965).

19M. Beyeler and D. Lazarus, Europhysics Conference on Atomic Transport in Solids and Liquids, Marstrand, Sweden, 1970 (unpublished).

20 L. W. Barr and J. N. Mundy, Diffusion in Body-Centered Cubic Metals (American Society for Metals, Metals Parks, Ohio, 1965), p. 171; J. N. Mundy, L. W. Barr, and F. A. Smith, Phil. Mag. 14, 785 (1966).

²¹A. P. Batra, Phys. Rev. <u>159</u>, 487 (1967).

22N. L. Peterson and S. J. Rothman, Phys. Rev. 163, 645 (1967).

23L. Chabildas and H. M. Gilder (unpublished).

²⁴A. D. LeClaire, Phil. Mag. <u>14</u>, 1271 (1966).

PHYSICAL REVIEW B

VOLUME 3, NUMBER 12

15 JUNE 1971

ERRATA

F-Center Gap Mode in Alkali Halides. A Molecular Model, R. S. Singh and S. S. Mitra [Phys. Rev. B 2, 1070 (1970)]. Dr. D. Bäuerle has brought to our attention certain misinterpretations of his results. The sentences starting from line 3 and ending in line 16 of the second paragraph of p. 1072 should accordingly be changed to "... For the F center in KI, a sharp band at 83 cm-1 has been observed in the far-infrared absorption. Clearly this should be compared with the calculated frequency of 81.2 cm $^{-1}$ (F_{1u}'). More recently Bäuerle and Hübner [Phys. Rev. B 2, 4252 (1970)] have reported the observation of \overline{a} gap mode due to an F center in KBr at 99.6 cm-1 (1.2 °K). This compares well with our calculated value of 92.6 cm-1 (300 °K) for this center. A resonant-band mode peaking around 62 cm⁻¹ has been reported^{6,14} for the KI:H- system."

Nonresonant Interband Faraday Rotation in Silicon, Cedric J. Gabriel [Phys. Rev. B 2, 1812 (1970)]. A printer's error was made in Eq. (3). The correct form of the equation is

$$2\gamma = \left[\xi(-) - \xi(+)\right] \left(\frac{d\xi(\pm)}{d\alpha}\right)^{-1}$$

High-Temperature Measurements of the Electron Hall Mobility in the Alkali Halides, C. H. Seager and David Emin [Phys. Rev. B 2, 3421 (1970)]. To correct for an error which has been discovered in the calibration procedure of the apparatus, the quoted Hall mobilities for all samples measured should be multiplied by a factor of 1.24 to obtain the proper values. This correction in no way affects the conclusion drawn by the authors that continuum polaron theories are inadequate in their predictions of the temperature dependence of the data.

Lattice-Dynamical Theory of the Diffusion Process. I. Isotope Effect in Cubic Metals, B. N. Narahari Achar [Phys. Rev. B 2, 3848 (1970)]. The 3N normal modes of the lattice perturbed by an isotopic impurity should be labeled by a single index f, rather than the symbols (\dot{q}, λ) . Accordingly, the symbols $(\overset{\star}{\mathbf{q}}, \lambda)$ should be replaced by f in the following: (i) Line above Eq. (10), p. 3850. (ii) Equations (10) and (11), p. 3850. (iii) Line 3 from top of the right-hand column, p. 3850.

Ionic Raman Effect. I. Scattering by Localized Vibration Modes, A. A. Maradudin and R. F. Wallis [Phys. Rev. B 2, 4294 (1970)]. An unfortunate numerical error occurred in plotting Figs. 1 and 2 on the basis of Eqs. (17) and (29), respectively. The scattering efficiencies shown in these figures are consequently too large by factors of 103-104. The corrected results, together with a demonstration that the ionic Raman effect for the system CaF2:H-dominates the electronic Raman effect when the incident light is in the infrared, are presented in a paper by L. B. Humphreys, A. A. Maradudin, and R. F. Wallis to appear in Physics of Impurity Centres in Crystals (Academy of Sciences of the Estonian S.S.R., Tartu, 1971).

Electromodulation of the Optical Properties of Thallium-Activated Potassium Bromide, U. Giorgianni, V. Grasso, and G. Saitta [Phys. Rev. B 2, 5007 (1970)]. Figures 1 and 2 should be interchanged,