

- ⁴ J. R. Ferraro, S. S. Mitra, and C. Postmus, *Phys. Rev. Letters* **18**, 455 (1967).
- ⁵ J. R. Ferraro, C. Postmus, S. S. Mitra, and C. J. Hoskins, *Appl. Opt.* **9**, 5 (1970).
- ⁶ J. R. Ferraro, S. S. Mitra, C. Postmus, C. Hoskins, and E. C. Siwiec, *Appl. Spectry.* **24**, 187 (1970).
- ⁷ J. Hiraishi and T. Shimanouchi, *Spectrochim. Acta* **22**, 1483 (1966).
- ⁸ Γ_T is the total number of translatory lattice modes.
- ⁹ H. Poulet, P. Delmore, and J. P. Mathieu, *Spectrochim. Acta* **20**, 1855 (1964).
- ¹⁰ A. Sabatini, L. Sacconi, and V. Schettino, *Inorg. Chem.* **3**, 1775 (1964).
- ¹¹ D. M. Adams and H. A. Gebbie, *Spectrochim. Acta* **19**, 925 (1963). D. M. Adams and D. C. Newton, *J. Chem. Soc.* **1969**, 2998.
- ¹² Γ_n is the total number of internal modes in the polyatomic unit MCl_4^{2-} .
- ¹³ H. G. Drickamer, R. W. Lynch, R. L. Clendenen, and E. A. Perez-Albuerno, *Advan. Solid State Phys.* **19**, 135 (1966).
- ¹⁴ J. Hiraishi, I. Nakagawa, and T. Shimanouchi, *Spectrochim. Acta* **20**, 819 (1964).