

57, 4401 (1972)], the tetragonal axis was aligned parallel to the direction of the temperature gradient in their Raman investigation. However, there is no evidence that this should always be true, and the present shear velocity data indicate that it is not.

<sup>10</sup>A. Bonilla, C. W. Garland, and N. E. Schumaker, *Acta Crystallogr. A* **26**, 156 (1970).

<sup>11</sup>V. Hovi, K. Heiskanen, and M. Varteras, *Ann. Acad. Sci. Fenn.* No. A144 (1964); V. Hovi, K. Paavola, and O. Urvaa, *Ann. Acad. Sci. Fenn.* No. A291 (1968).

<sup>12</sup>C. W. Garland and R. A. Young, *J. Chem. Phys.* **48**, 146 (1968).

<sup>13</sup>T. S. Velichkina, O. N. Golubeva, O. A. Shustina, and I. A. Yakovlev, *Bull. Acad. Sci. USSR Phys. Ser.* **35**, 932 (1971).

<sup>14</sup>K. Kawasaki, *Int. J. Magn.* **1**, 171 (1971).

<sup>15</sup>K. Kawasaki, *Phys. Rev. A* **1**, 1750 (1970).

<sup>16</sup>B. I. Halperin and P. C. Hohenberg, *Phys. Rev.* **177**, 952 (1969).

<sup>17</sup>G. E. Laramore and L. P. Kadanoff, *Phys. Rev.* **187**, 619 (1969).

<sup>18</sup>E. Riedel and F. Wegner, *Phys. Rev. Lett.* **24**, 730 (1970); E. K. Riedel, *J. Appl. Phys.* **42**, 1383 (1971).