

- \*Work supported in part by the NSF.
- <sup>1</sup>B. B. Weiner and C. W. Garland, J. Chem. Phys. 56, 155 (1972); see also C. W. Garland and B. B. Weiner, Phys. Rev. B 3, 1634 (1971).
- <sup>2</sup>(a) C. W. Garland, D. E. Bruins, and T. J. Greytak, Phys. Rev. B 12, 2759 (1975); (b) D. E. Bruins and C. W. Garland, J. Chem. Phys. 63, 4139 (1975).
- <sup>3</sup>C. C. Stephenson (private communication of unpublished data).
- <sup>4</sup>H. Chihara and M. Nakamura, Bull. Chem. Soc. Jpn. 45, 133 (1972).
- <sup>5</sup>E. B. Amitin, Yu. A. Kovalevskaya, E. G. Lebedeva, and I. E. Paukov, Fiz. Tverd. Tela 17, 2549 (1975) [Sov. Phys.-Solid State 17, 1699 (1976)].
- <sup>6</sup>A. V. Voronel and S. R. Garber, Zh. Eksp. Teor. Phys. 52, 1464 (1967) [Sov. Phys.-JETP 25, 970 (1967)].
- <sup>7</sup>P. Schwartz, Phys. Rev. B 4, 920 (1971); and Ph.D. thesis (Univ. of Illinois, 1969) (unpublished).
- <sup>8</sup>S. A. Zlunitsyn, Zh. Eksp. Teor. Fiz. 8, 724 (1938).
- <sup>9</sup>E. B. Amitin, Yu. A. Kovalevskaya, and I. E. Paukov, Fiz. Tverd. Tela 14, 3483 (1972) [Sov. Phys.-Solid State 14, 2902 (1973)].
- <sup>10</sup>N. J. Trappeniers and Th. J. van der Molen, Physica 32, 1161 (1966); Th. J. van der Molen, Ph.D. thesis (Univ. of Amsterdam, 1966) (unpublished).
- <sup>11</sup>J. D. Baloga and C. W. Garland, Rev. Sci. Instrum. (to be published).
- <sup>12</sup>R. Renard and C. W. Garland, J. Chem. Phys. 45, 763 (1966).
- <sup>13</sup>C. W. F. T. Pistorius, J. Chem. Phys. 50, 1436 (1969).
- <sup>14</sup>W. Mandema and N. J. Trappeniers, Physica 76, 123 (1974).
- <sup>15</sup>W. Bausch and W. Waidelich, Phys. Lett. A 30, 190 (1969).
- <sup>16</sup>S. S. Sharma, Proc. Indian Acad. Sci. A 31, 339 (1950).
- <sup>17</sup>No measurements were made on this sample at 1 atm, and these *r* values should not be directly compared with that for sample *B*.
- <sup>18</sup>An additional confirmation of this reversible reduction of the peak *C<sub>p</sub>* value with pressure was obtained from a series of measurements on a third ND<sub>4</sub>Cl sample. Data were taken only in the immediate vicinity of the transition for isobars at 760, 1100, 1500, and 2730 bar. A smooth monotonic decrease in peak *C<sub>p</sub>* values was observed with increasing pressure, and the 760 bar value was shown to be reproducible after pressure cycling. Our thanks to G. Kasting for carrying out this investigation.
- <sup>19</sup>D. G. Thomas and L. A. K. Staveley, J. Chem. Soc. Lond. A 1951, 1420 (1951).
- <sup>20</sup>G. E. Fredericks, Phys. Rev. B 4, 911 (1971).
- <sup>21</sup>T. S. Chang, G. F. Tuthill, and H. E. Stanley, Phys. Rev. B 9, 4882 (1974).
- <sup>22</sup>W. B. Yelon and D. E. Cox, Solid State Commun. 11, 1011 (1972); W. B. Yelon, D. E. Cox, P. J. Kortman, and W. B. Daniels, Phys. Rev. B 9, 4843 (1974).
- <sup>23</sup>D. R. McKenzie and R. S. Seymour, J. Phys. C 8, 1071 (1975).