

substituted YFe garnet system. The agreement between the two methods is quite good, but is actually even better than it looks if we take the 0°K moments that the other authors obtained for their specimens (see Ref.<sup>27</sup>).




Fig. 6. Fraction,  $f_t$ , of  $Ga^{3+}$  and  $Al^{3+}$  ions in tetrahedral sites in  $Y_3Fe_{5-x}Ga_xO_{12}$  and  $Y_3Fe_{5-x}Al_xO_{12}$ , respectively (from Ref.<sup>27</sup>)




Fig. 7. Lattice constant vs  $x$  (from Ref.<sup>27</sup>)

For  $x > 1.50$ , the distribution appears to be sensitive to specimen heat treatment and this may be part of the reason for different 0°K moments obtained by different investigators. The importance of care in specimen preparation cannot be overestimated. In many systems the lattice-constant measurements can give an indication of the

specimen quality. Smooth curves should be obtained for lattice constant and moment, at a fixed temperature, vs composition. Examples from the  $Y_3Fe_{5-x}Ga_xO_{12}$  system are given in Figs. 7 and 8.

Fig. 6 shows that for most of the composition range the  $Ga^{3+}$  ions have a greater preference for tetrahedral sites ( $f_t =$  fraction of  $Ga^{3+}$  or  $Al^{3+}$  ions in tetrahedral sites) than  $Al^{3+}$  ions to  $x \approx 2.75$ . For  $x > 2.75$ , the curves may coincide or cross. Unfortunately, we cannot




Fig. 8. Spontaneous moment at 0°K vs  $x$  (from Ref.<sup>27</sup>). (See Ref.<sup>27</sup> for pertinent references)

learn from static magnetic measurements anything about ionic distributions in specimens with  $x$  greater than 3.00. Furthermore, it is unlikely that the accuracy desired is attainable by diffraction techniques.

In regard to the diffraction techniques, a paper by FISCHER *et al.*<sup>101</sup> purports to give the distributions in the systems by both x-ray and

<sup>101</sup> P. FISCHER, W. HÄLG, E. STOLL and A. SEGMÜLLER, X-ray and neutron diffraction study of substitutional disorder in yttrium-iron-gallium garnets. *Acta Crystallogr.* 21 (1966) 765-769.