

sion a Mg-poor calcite I could be exsolved. In either case, there would be a resulting confusion as to the original composition of the primary phase. The present data indicate that such an effect is not of real concern, because the differential solubility is so slight that the geologic thermometry would not be affected. It is a consequence of the small solubility differential that the field of calcite II remains at high temperatures and pressures and that calcite II as a primary phase is probably not often encountered in ordinary metamorphic conditions. For example, if the temperature of the Crestmore, California, carbonates as interpreted by Carpenter (1967) is correct, calcite II would have been the primary phase if the pressure exceeded approximately 5 kb. This represents a depth of burial in excess of 15 km, which, although perhaps not impossible, is unlikely.

The presence of MgCO<sub>3</sub> increases the field of stability of calcite with respect to aragonite. The effect is rather small at temperatures up to approximately 450°C, but at higher temperatures the difference between the transition curves for pure CaCO<sub>3</sub> and for the MgCO<sub>3</sub>-saturated system becomes large. It is unlikely, however, that aragonite is produced in nature in the higher grades of metamorphism, as the necessary pressures are unrealistically high. The presence of MgCO<sub>3</sub> in the system makes it all the more unlikely that aragonite would be formed under equilibrium conditions. Metamorphic aragonite is known from several low-grade metamorphic terranes (Coleman and Lee, 1962). These rocks probably never reached temperatures higher than about 250°C, at which point the pure calcite-aragonite and MgCO<sub>3</sub>-saturated calcite-aragonite curves are separated by but a few hundred bars. Therefore, the presence of MgCO<sub>3</sub> in rocks probably has little effect on a geobarometer based on the equilibrium calcite-aragonite transition.

#### ACKNOWLEDGMENTS

We are indebted to Professors P. J. Wyllie and John C. Jamieson for a critical reading of the manuscript. This work was supported by NSF Grants No. GP-2695 and GA-573 and by PRF Grant No. 3204-C2 and ARPA Grant SD-89.

#### REFERENCES

- Bell, P. M., and England, J. L., 1964, High pressure differential thermal analysis of a fast reaction with CaCO<sub>3</sub>: Carnegie Inst. Washington Year Book 63, p. 176-178.
- Boeke, H. E., 1912, Die Schmelzerscheinungen und die umkehrbare Umwandlung des Calcium-Carbonats: Neues Jahrb. Mineralogie Geologie, v. 1, p. 91-212.
- Boettcher, A. L., and Wyllie, P. J., 1967, Revision of the calcite-aragonite transition, with the location of a triple point between calcite I, calcite II, and aragonite: Nature, v. 213, p. 792-793.
- , 1968, The calcite-aragonite transition measured in the system CaO-H<sub>2</sub>O-CO<sub>2</sub>: Jour. Geology v. 76, p. 314-330.
- Boyd, F. R., and England, J. L., 1960, Apparatus for phase equilibrium measurements at pressures up to 50 kilobars and temperatures up to 1750°C: Jour. Geophys. Research, v. 65, p. 741-748.
- Bridgman, P. W., 1939, The high pressure behavior of miscellaneous materials: Am. Jour. Sci., v. 237, p. 7-18.
- , 1952, The resistance of 72 elements, alloys and compounds to 100,000 KG/CM<sup>2</sup>: Am. Acad. Arts and Sci. Proc., v. 81, p. 165-250.

- Carpenter, A. B., 1967, Mineralogy and petrology of the system  $\text{CaO-MgO-CO}_2\text{-H}_2\text{O}$  at Crestmore, California: *Am. Mineralogist*, v. 52, p. 1341-1365.
- Clark, S. P. Jr., 1957, A note on calcite-aragonite equilibrium: *Am. Mineralogist*, v. 42, p. 564-566.
- Cohen, L. H., and Klement, W. K. Jr., 1967, High-low quartz inversion. Determination to 35 kilobars: *Jour. Geophys. Research*, v. 72, p. 4245-4251.
- Coleman, R. G., and Lee, D. E., 1962, Metamorphic aragonite in the glaucophane schists of Cazadero, California: *Am. Jour. Sci.*, v. 260, p. 577-595.
- Crawford, W. A., and Fyfe, W. S., 1964, Calcite-aragonite equilibrium at 100°C: *Science*, v. 144, p. 1569-1570.
- Dandekar, D., ms, 1967, Variation in elastic constants of calcite with pressure: Ph.D. thesis, Dept. Geophys. Sci., Univ. Chicago.
- Eitel, W., and Skaliks, W., 1929, Über einige Doppelcarbonate der Alkalien und Erdalkalien: *Zeitschr. Anorg. Allg. Chemie*, v. 183, p. 268-286.
- Fyfe, W. S., Turner, F. J., and Verhoogen, J., 1958, Metamorphic reactions and metamorphic facies: *Geol. Soc. America Mem.* 73, 259 p.
- Goldsmith, J. R., 1959, Some aspects of the geochemistry of carbonates, in Abelson, P. H., ed., *Researches in geochemistry*: New York, John Wiley & Sons, p. 336-358.
- Goldsmith, J. R., and Graf, D. L., 1960, Subsolidus relations in the system  $\text{CaCO}_3\text{-MgCO}_3\text{-MnCO}_3$ : *Jour. Geology*, v. 68, p. 324-335.
- Goldsmith, J. R., Graf, D. L., and Heard, H. C., 1961, Lattice constants of the calcium-magnesium carbonates: *Am. Mineralogist*, v. 46, p. 453-457.
- Goldsmith, J. R., Graf, D. L., Witters, J., and Northrop, D. A., 1962, Studies in the system  $\text{CaCO}_3\text{-MgCO}_3\text{-FeCO}_3$ : 1. Phase relations; 2. A method for major element spectrochemical analysis; 3. Compositions of some ferroan dolomites: *Jour. Geology*, v. 70, p. 659-688.
- Goldsmith, J. R., and Heard, H. C., 1961, Subsolidus phase relations in the system  $\text{CaCO}_3\text{-MgCO}_3$ : *Jour. Geology*, v. 69, p. 45-74.
- Graf, D. L., and Goldsmith, J. R., 1955, Dolomite-magnesian calcite relations at elevated temperatures and  $\text{CO}_2$  pressures: *Geochim. et Cosmochim. Acta*, v. 7, p. 109-128.
- \_\_\_\_\_ 1958, The solid solubility of  $\text{MgCO}_3$  in  $\text{CaCO}_3$ : A revision: *Geochim. et Cosmochim. Acta*, v. 13, p. 218-219.
- Guggenheim, E. A., 1952, *Mixtures*: Oxford, Clarendon Press, 270 p.
- Hanneman, R. E., and Strong, H. M., 1965, Pressure dependence of electromotive force of thermocouples to 1300°C and 50 kilobars: *Am. Soc. Mining Engineers Ann. Winter Mtg.*, New York 1964, ASMR Tech. Paper 64-WA/PT-21.
- Harker, R. I., and Tuttle, O. F., 1955, Studies in the system  $\text{CaO-MgO-CO}_2$ : Pt. 2: Limits of solid solution along the binary join,  $\text{CaCO}_3\text{-MgCO}_3$ : *Am. Jour. Sci.*, v. 253, p. 274-282.
- Jamieson, J. C., 1957, Introductory studies of high pressure polymorphism to 24,000 bars by X-ray diffraction, with some comments on calcite II: *Jour. Geology*, v. 65, p. 334-344.
- Kennedy, G. C., and LaMori, P. N., 1962, The pressures of some solid-solid transitions: *Jour. Geophys. Research*, v. 67, p. 851-856.
- Kitahara, S., and Kennedy, G. C., 1964, The quartz-coesite transition: *Jour. Geophys. Research*, v. 69, p. 5395-5400.
- MacDonald, G. J. F., 1956, Experimental determination of calcite-aragonite relations at elevated temperatures and pressures: *Am. Mineralogist*, v. 41, p. 744-756.
- Robie, R. A., ms, 1957, Thermodynamic properties of  $\text{CaMg}(\text{CO}_3)_2$ ,  $\text{Mg}_3\text{Si}_4\text{O}_{10}(\text{OH})_2$  and  $\text{Ca}_2\text{Mg}_5\text{Si}_8\text{O}_{22}(\text{OH})_2$  between 12 and 300 degrees Kelvin: Ph.D. thesis, Dept. Geology, Univ. Chicago.
- Tuttle, O. F., 1949, Two pressure vessels for silicate-water studies: *Geol. Soc. America Bull.* v. 60, p. 1727-1729.