

- to 50 kilobars and temperatures up to 1750°C, *J. Geophys. Res.*, *65*, 741-748, 1960b.
- Boyd, F. R., and J. L. England, Effect of pressure on the melting point of diopside, CaMgSi₂O₆, and albite, NaAlSi₃O₈, in the range up to 50 kilobars, *J. Geophys. Res.*, *68*, 311-323, 1963.
- Khitarov, N. I., New experimental work in the field of deep-seated processes, *Geochemistry USSR, English Transl.*, no. 6, 578-581, 1964.
- Kitahara, S., and G. C. Kennedy, The quartz-coesite transition, *J. Geophys. Res.*, *69*, 5395-5400, 1964.
- Klement, W., Jr., A. Jayaraman, and G. C. Kennedy, Transformation in mercury at high pressures, *Phys. Rev.*, *131*, 1-6, 1963.
- Newton, R. C., The thermal stability of zoisite, *J. Geol.*, *73*, 431-441, 1965.
- Ringwood, A. E., and D. H. Green, Experimental investigations bearing on the nature of the Mohorovicic discontinuity, *Nature*, *201*, 566-567, 1964.

(Manuscript received February 18, 1966;
revised April 1, 1966.)

Glassy Spherule

Most students of cosmic magnetic opaque particles tallic, partly because of and partly because it is silicate particles from terr [Wright *et al.*, 1963; Hod, ever, we could hypothesize mic dust particles that are similar in compos.

In the precipitate of m December 20, 1964, near S glassy spherules were found scope. One of the typical shown in Figure 1. In sh with a tail-like protrusion face is bright, clear, and and no fissures are observed transparent body with no index is 1.505. It is not m composition is (in weight Al₂O₃, 12.5; FeO (total), 3.0; Na₂O, 1.5; K₂O, 2.5. A small, it was analyzed by method using the electron

Fig. 1. The typical 'tear