

- a basic parent. Thus  
ma the mass ratio of  
ia produced is ap-  
models involving  
pressure phases (e.g.  
t) the mass ratio of  
ould be two or more,
- [15] P. W. Lipman, Mineralogy and paragenesis of amphibole  
from Gibson Peak pluton, Northern California, Am.  
Min. 49 (1964) 1321.
- [16] W. A. Deer, The composition and paragenesis of the  
hornblendes of the Glen Tilt Complex, Perthshire, Min.  
Mag. 25 (1938) 56.
- [17] H. H. Hess, History of ocean basins, in: A. E. J. Engel,  
H. L. James and B. F. Leonard (eds.), Petrologic  
Studies – Buddington Vol. Geol. Soc. Am., N.Y., 599.
- [18] T. H. Green, D. H. Green and A. E. Ringwood, The ori-  
gin of high-alumina basalts and their relationships to  
quartz tholeiites and alkali basalts, Earth Planet. Sci.  
Letters 2 (1967) 41.
- [19] D. H. Green and A. E. Ringwood, Genesis of basaltic  
magmas, Contr. Mineral. Petrol. 15 (1967) 103.
- [20] A. E. Ringwood and D. H. Green, An experimental in-  
vestigation of the gabbro-eclogite transformation and  
some geophysical implications, Tectonophys. 3 (1966)  
383.