

down" metamorphism. Brothers ruled out deep tectonic burial on grounds of the passage downward into "lower pressure" rocks. He ruled out tectonic overpressure on grounds of textural evidence for static recrystallization. Brothers proposed a new hypothesis of gas overpressure by vapors trapped below an impermeable ultramafic caprock. But the relationships described by him are also compatible with my model of development of unusual chemical conditions in pore fluids near ultramafic bodies undergoing serpentinization.

In summary, the spatial association of blueschists with serpentinites is firmly established on a worldwide basis. In regard to the possible genetic significance of the association, I urge the interested reader to take neither my word nor Ernst's as the final authority. Rather, he should read the available literature and let the evidence speak for itself.

REFERENCES

- Bailey, E. H., Blake, M. C., Jr., and Jones, D. L., 1970, On-land Mesozoic oceanic crust in California Coast Ranges, in Geological Survey Research 1970: U.S. Geol. Survey Prof. Paper 700-C, p. C70-C81.
- Black, P. M., 1970a, Ferroglaucophane from New Caledonia: *Am. Mineralogist*, v. 55, p. 508-511.
- 1970b, P2 omphacite intermediate in composition between jadeite and hedenbergite, from metamorphosed acid volcanics, Bouehndep, New Caledonia: *Am. Mineralogist*, v. 55, p. 512-514.
- Blake, M. C., Jr., Irwin, W. P., and Coleman, R. G., 1969, Blueschist-facies metamorphism related to regional thrust faulting: *Tectonophysics*, v. 8, p. 237-246.
- Brothers, R. N., 1970, Lawsonite-albite schists from northernmost New Caledonia: *Contr. Mineralogy and Petrology*, v. 25, p. 185-202.
- Brouwer, H. A., and Egeler, C. G., 1952, The glaucophane facies metamorphism in the schistes lustrés nappe of Corsica: *Koninkl. Nederlandse Akad. Wetensch., Afdeling Natuurk.*, (Tweede Reeks), pt. 48, no. 3, 71 p.
- Chamberlin, J. A., McLeod, C. R., Traill, R. J., and Lachance, G. R., 1965, Native metals in the Muskox intrusion: *Canadian Jour. Earth Sci.*, v. 2, p. 188-215.
- Christensen, N. I., 1971, Fabric, seismic anisotropy, and tectonic history of the Twin Sisters dunite: *Geol. Soc. America Bull.*, in press.
- Cogulu, E., 1967, Etude pétrographique de la région de Mihalicik (Turquie): *Suisse Mineralogic Petrographie Bull.*, v. 47, p. 683-824.
- Coleman, R. G., 1967, Glaucophane schists from California and New Caledonia: *Tectonophysics*, v. 4, p. 479-498.
- 1971, Plate tectonic emplacement of upper mantle peridotites along continental edges: *Jour. Geophys. Research*, v. 76, p. 1212-1222.
- Coombs, D. S., and Landis, C. A., 1966, Metamorphic belts and orogenesis in New Zealand [abs.], in *Age and Nature of the Circum-Pacific Orogenesis: Pacific Sci. Cong.*, 11th, Tokyo, Proc., v. 4, p. 13.
- Daniels, W. B., and Skoultchi, A. I., 1966, Optical observations of the f.c.c.s.c. transformation in single crystals of RBI at high pressures: *Jour. Phys. Chem. Solids*, v. 27, p. 1247-1250.
- Davies, H. L., ms, 1969, Peridotite-gabbro-basalt complex in eastern Papua: An overthrust plate of oceanic mantle and crust: Ph.D. thesis, Stanford Univ.
- deRoever, W. P., 1947, Igneous and metamorphic rocks in eastern Central Celebes, in *Geological Explorations in the Island of Celebes under the Leadership of H. A. Brouwer: Amsterdam, North Holland Pub. Co.*, p. 65-173.
- Dobretsov, N. L., and Kuroda, I., 1970, Geologic law characterizing glaucophane metamorphism in northwestern part of the folded frame of Pacific Ocean: *Internat. Geology Rev.*, v. 12, p. 1389-1407.
- Ernst, W. G., 1957, Annual Report of the Director of the Geophysical Laboratory, 1956-1957: *Carnegie Inst. Washington Year Book* 56, p. 228.
- 1971, Do mineral parageneses reflect unusually high-pressure conditions of Franciscan metamorphism?: *Am. Jour. Sci.*, v. 270, p. 81-108.

- Essene, E. J., ms, 1961, Petrogenesis of Franciscan metamorphic rocks: Ph.D. thesis, Univ. California, Berkeley.
- Gresens, R. L., 1969, Blueschist alteration during serpentinization: *Contr. Mineralogy and Petrology*, v. 24, p. 93-113.
- Joplin, G. A., 1968, *A Petrography of Australian Metamorphic Rocks*: New York, Am. Elsevier Pub. Co.
- Martin, R. F., 1969, The hydrothermal synthesis of low albite: *Contr. Mineralogy and Petrology*, v. 23, p. 323-339.
- Milton, C., and Eugster, H. P., 1959, Mineral assemblages of the Green River formation, in Abelson, P. H., ed., *Researches in Geochemistry*: New York, John Wiley and Sons, Inc.
- Miyashiro, Akiho, 1961, Evolution of metamorphic belts: *Jour. Petrology*, v. 2, p. 277-311.
- Quodling, F. M., 1964, On traces of native iron at Port Macquarie, New South Wales: *Royal Soc. New South Wales Jour. and Proc.*, v. 97, p. 81-82.
- Shagam, R., 1960, Geology of central Aragua, Venezuela: *Geol. Soc. America Bull.*, v. 71, p. 249-302.
- Taylor, H. P., Jr., and Coleman, R. G., 1968, O^{18}/O^{16} ratios of coexisting minerals in glaucophane-bearing metamorphic rocks: *Geol. Soc. America Bull.*, v. 79, p. 1727-1756.
- Thayer, T. P., 1966, Serpentinization considered as a constant-volume metasomatic process: *Am. Mineralogist*, v. 51, p. 685-710.
- Williams, H., McBirney, A. R., and Dengo, G., 1964, Geologic reconnaissance of southeastern Guatemala: *Univ. California Pub. Geol. Sci.*, v. 50, p. 1-62.
- Yegorov, A. Y., Dobretsov, N. L., Yegorova, M. G., and Podzorova, D. I., 1967, Glaucophane schists of Sakhalin Island (Eng. translation): *Akad. Nauk SSSR Doklady*, v. 175, p. 169-171.

REPLY

W. G. ERNST

Department of Geology and Institute of
Geophysics and Planetary Physics,
University of California, Los Angeles, California 90024

Progressive, areally distinct, at least in part contemporaneous metamorphic sequences developed in *in situ* Franciscan rocks have been elucidated by many workers. (For documentation of this and other aspects of the problem, see references cited in Ernst, 1971, and Gresens, 1971.) In quartz and layer silicate-bearing metagraywackes, the most typical paragenesis seems to be: (A) laumontite + albite \pm calcite; (B) pumpellyite + albite \pm calcite; (C) lawsonite + albite \pm calcite or aragonite; and (D) lawsonite + jadeitic pyroxene \pm aragonite. Mafic metavolcanics exhibit a corresponding progression from feebly recrystallized greenstones to blueschists. The available thermochemical data and numerous experimental phase equilibrium studies are consistent with the observed mineralogic changes and suggest that relatively high pressures attended metamorphism. Combined with oxygen isotope geothermometry, it would appear that aragonite- and jadeitic pyroxene-bearing metagraywackes and associated metavolcanics must have crystallized at pressures exceeding 8 kb at temperatures on the order of 150 to 300°C. (Some investigators, while accepting these high pressures, have postulated substantial stress increment as the means whereby high pressures could have accompanied such low temperatures of metamorphism. However,