

TABLE VII
CONSTANTS^a IN EQ. (4) FOR MOLAR VOLUMES OF FLUID ALONG THE MELTING CURVE

Fluid	a'	b'	c'	d'	P_m range, kg/cm ²	rms dev., cm ³ /mol
He ⁴ II	0	-0.17145	1	27.570	26-30	0.0006
He ⁴ I	14.854	48.5273	-0.107253	-10.0712	35-3555	0.0097
He ³	1.075	51.1102	-0.161532	-3.2482	50-3555	0.0137

^a Pressure units in kg/cm² and volume units in cm³/mol.

2. Thermal expansion and compressibility of the fluid

The thermal expansion coefficient of fluid He³ along the melting curve exhibits a maximum in the vicinity of the triple point, as shown in Figs. 3 and 4. The maximum is broad compared to that for He⁴ and is less than one-half as large. In general, one expects α to increase with T and decrease with P ; however, along the melting curve the "normal" behavior of α_f increasing with decreasing P_m and T_m indicates that P_m changes overcome T_m changes. For He⁴ the maximum in α_f appears to be a direct consequence of the λ -transition. In He³ the nuclear spin part of α_f becomes more negative at lower T , according to Goldstein (25), and it apparently overcomes the "normal" behavior of the nonspin part of α_f .

From values of α_f and β_f in Fig. 4, it is possible to compute the following thermodynamic quantities for fluid He³ along the melting curve:

$$(\partial P / \partial T)_V = \alpha_f / \beta_f; \quad (5)$$

and

$$(C_P - C_V) = TV_f \alpha_f^2 / \beta_f. \quad (6)$$

These quantities are shown as the curves in Fig. 10. Neither curve exhibits a maximum over the range studied. The plot of $(C_P - C_V)$ versus P_m is linear below 180 kg/cm² and extrapolates to zero at $P_m = 47$ kg/cm². This extrapolation gives a good determination of the point where α_f goes through zero on the melting curve.

The pressure-temperature locus of $\alpha_f = 0$ in the fluid domain is shown in Figs. 5 and 9. For completeness, the point of Taylor and Kerr (26) on the vaporization curve has been included.³ The points represented by open circles were obtained by extrapolation to zero of a series of α_f values measured at constant pressure and various temperatures. This could be done reliably because the slopes were quite constant. Extrapolations were made below about 1.4°K, the

³ Lee *et al.* (27) also reported a density maximum at approximately 0.5°K, presumably at saturation.