

- ZADWORSKI, J. AND S. BRETSZKAPFER (1933a) Heterogeneous reactions of the type $A \text{ solid} + B \text{ gas} = C \text{ solid}$: I. Deviations from constancy of the equilibrium pressure; apparatus equilibrium and its significance. *Z. Physik. Chem.*, **35**, 60-78.
- _____, and _____ (1933b) Heterogeneous reactions of the type $A \text{ solid} + B \text{ gas} = C \text{ solid}$.
- II. Kinetics of carbonate formation and decomposition. *Z. Physik. Chem.*, **35**, 79-96.
- _____, and _____ (1938a) Heterogeneous reactions of the type $A \text{ solid} + B \text{ gas} = C \text{ solid}$.
- III. Experimental determination of the elementary reactions in the combined process.
- Z. Physik. Chem.*, **40**, 158-182.
- _____, and _____ (1938b) Mechanism of reaction of the type $\text{solid} + \text{gas} = \text{solid}$. *Trans. Faraday Soc.*, **34**, 981-939.
- ZARSHNEVSKAYA, N. G. (1964) Origin of the gases in rocks of the Khibina apatite deposits. *Dokl. Akad. Nauk SSSR*, **154**, 118-120. [*Chem. Abstr.* **60**, 9039 [1964]; Engl. transl. *Dokl. Acad. Sci. USSR, Earth Sci. Sec.* **154**, 128-131].
- Manuscript received, July 27, 1968; accepted for publication, May 1, 1969.*

Chemical and modal analysis of ultramafic body in Noronha, and an increase in volume

Although many petro-

are available (Thayer, 1966, 1967; Himm-

sufficient to determine

accompany progressive

the inability to resolve

chemical and mineralogical

the degree of serpentinization

Such data are present in

dunite samples from the

The Webster-Addie mafic

bodies in the Apennine

Sylvania. It is elliptical

outward, and varies in

A complete description

in preparation (Madison

chiefly of dunite, with

serpentine). Most of the

In order to limit the

stearitization, only petro-

Twenty-two parts

investigation.

Fe, Mg, Ca, and Ni