

- ZADOWSKI, J. AND S. BRETSZNAJDER (1933a) Heterogeneous reactions of the type A solid + B gas = C solid: I. Deviations from constancy of the equilibrium pressure; apparent equilibrium and its significance. *Z. Physik. Chem.*, **35**, 60-78.
- , and ——— (1933b) Heterogeneous reactions of the type A solid + B gas = C solid: II. Kinetics of carbonate formation and decomposition. *Z. Physik. Chem.*, **35**, 79-96.
- , and ——— (1938a) Heterogeneous reactions of the type A solid + B gas = C solid: III. Experimental determination of the elementary reactions in the combined process. *Z. Physik. Chem.*, **40**, 158-182.
- , and ——— (1938b) Mechanism of reaction of the type solid = solid + gas: *Trans. Faraday Soc.*, **34**, 951-959.
- ZAKRSHEVSKAYA, N. G. (1964) Origin of the gases in rocks of the Khibina apatite deposits. *Dokl. Akad. Nauk SSSR*, **154**, 118-120. [*Chem. Abstr.* **60**, 9039 [1964]; Engl. transl. *Dokl. Acad. Sci. USSR, Earth Sci. Sec.* **154**, 128-131].

Manuscript received, July 27, 1968; accepted for publication, May 1, 1969.

COMPOSITIONAL
PANYING PROG
DUNITES FROM T
BOD

KENT C. CONDIE, I
Univers

JAMES A. MADISON, I
Univers

Chemical and modal analy
Addie ultramafic body in Nor
Na, and an increase in volume

Although many petro
are available (Thayer, I
Page, 1966, 1967; Him
sufficient to determine
accompany progressive
the inability to resolve
chemical and mineralog
the degree of serpentin
Such data are present
dunite samples from th
lina.

The Webster-Addie
mafic bodies in the Ap
northwestern part of J
Sylva. It is elliptical
outward, and varies in
A complete description
in preparation (Madis
chiefly of dunite, with
sterite). Most of the r
In order to limit the
steatitization, only pe
investigation.

Twenty-two partic
Fe, Mg, Ca, and Ni